

11.0 ECOLOGICAL RISK CONCLUSIONS

Risk estimates in this BERA were calculated following CERCLA guidance (EPA 1997, 1998) and EPA's Problem Formulation (Attachment 2). The conclusions of the BERA, along with those of the BHHRA (Appendix F of the draft final RI (Integral et al. 2011)), are intended to provide information to risk managers on potentially unacceptable risks predicted under current conditions of the Study Area, as well as information on possible future approaches for protecting human health and the environment.

The BERA's risk conclusions are provided at the end of the risk assessment for each receptor group:

- In Section 6.7 for benthic community assessment endpoints
- In Section 7.6 for fish assessment endpoints
- In Section 8.3 for avian and mammalian assessment endpoints
- In Section 9.3 for the amphibian assessment endpoint
- In Section 10.3 for the aquatic plant assessment endpoint

Consistent with ERAGs (EPA 1997) the foregoing risk conclusions identified the receptor-COPC pairs that, given the magnitude and extent of risk, are reasonably likely to result in adverse effects on the assessment endpoints selected to represent the valued ecological attributes of the Study Area. Section 11 does not recapitulate the analyses that went into drawing the risk conclusions. For that level of detail the reader is referred back to the aforementioned risk conclusion sections. The remainder of Section 11 is organized as follows:

- Section 11.1 presents a summary by receptor group and LOE of the 89⁺ 102 ecological COPCs identified as posing potentially unacceptable risk in this BERA based on $HQ \geq 1$ for at least one receptor-LOE combination.
- Section 11.2 identifies COPCs identified as posing potentially unacceptable risks for ecological receptors in the Study Area that occur at concentrations similar to the sediment and surface water background levels defined in Section 7.0 of the draft final RI (Integral et al. 2011) or to tissue concentrations in four fish receptor species (i.e., juvenile Chinook salmon, brown bullhead, smallmouth bass, and lamprey ammocoetes) collected from the upriver reach of the Willamette River (RM 15.3 to RM 28.4).
- Section 11.3 combines the risk conclusions across all ecological receptor groups to provide a general overview of ecological risks and to identify the

Comment [A1]: Footnote 1 was pointed out to be in error by EPA in our draft final BERA comments (Comment 132). TPH is a CERCLA contaminant. The TPH in sediment TRVs were given to LWG on April 11, 2008, including the methodology of their derivation and a table with the numeric values corresponding to Oregon's aromatic and aliphatic TPH fraction definitions. Section 5.2 of the BERA discusses these TPH TRVs, which should also have been used in the BERA in the absence of any modified TPH TRVs.

⁺ Ninety-one contaminants have $HQs \geq 1$. Because petroleum compounds are not CERCLA contaminants, gasoline-range hydrocarbons and diesel-range hydrocarbons have been excluded from the final count even though they may be contributing to potentially unacceptable risk.

receptor-COPC pairs that, given the magnitude and extent of risk, are reasonably likely to result in adverse effects on the assessment endpoints.

Risk management recommendations from the LWG risk assessors to EPA risk managers, based on the results of the BERA, are presented in Section 12.

11.1 SUMMARY OF POTENTIALLY UNACCEPTABLE RISKS

Consistent with EPA Superfund ERA guidance (EPA 1997, 1998), potentially unacceptable risks were identified through an iterative process of analyzing the exposure and effects data for the various chemicals and ecological receptors, with increasing realism at each step in the process. For most receptors, several LOEs were evaluated (Section 3.3). For each LOE, risk characterization began with the SLERA (Section 5) and progressed iteratively through the final step in the risk characterization. Throughout the process, chemical-receptor pairs that showed the potential for adverse effects were further analyzed and those that did not were screened out. The final step in the process reflects the most realistic risk estimates. Potentially unacceptable risks were identified for each receptor-LOE-COPC combination based on the final step in the risk characterization.

Exposure data in the final step of the risk analysis were evaluated at the scale over which the receptors are likely to be exposed and, where pertinent, the variety of potentially contaminated prey the receptor may consume. For the least mobile receptors (e.g., benthic macroinvertebrates, sculpin, aquatic plants), exposure areas are no larger than the immediate area where samples were collected; for the most mobile receptors (e.g., white sturgeon, largescale sucker), the exposure areas encompass the entire Study Area. For moderately mobile receptors (e.g., smallmouth bass, mink) the Study Area is divided into several exposure areas each 1 to 3 miles long.

For all LOEs except sediment, numerical risk estimates were calculated as HQs (Equation 6-1). HQs were calculated separately for each receptor-LOE-COPC combination for each exposure area. Receptor-LOE-COPC combinations resulting in $HQ \geq 1$ in the final step of the risk characterization in any exposure area were identified as posing potentially unacceptable risk. For the sediment LOE, a location was identified as posing potentially unacceptable risk to benthic invertebrates if the sediment was toxic or predicted to be toxic based on a sediment COPC concentration that exceeded a site-specific SQV.

Those chemicals for which exposure or effects data were insufficient to evaluate the risk were also identified as posing potentially unacceptable risk, although risk is unknown. [Potential risks from chemicals without TRVs cannot be quantified, and are an uncertainty in the BERA that may underestimate risks.](#) Risk to benthic organisms, including clams and crayfish, could not be evaluated for 78 sediment COIs because either no relationship between sediment contaminants and toxicity was apparent in the site-specific dataset or too few data points were available to discern a relationship (Table 6-6 summarizes the selection of chemicals for evaluation of site-specific toxicity). Other contaminants that could not be evaluated for their contribution to benthic community risks include 27 tissue

COIs (Table 6-28), 19 surface water COIs (Table 6-35), and 16 TZW COIs (Table 6-42). Risk to fish from a number of COIs could not be evaluated: 17 tissue-residue COIs (Table 7-13), 11 dietary COIs (Table 7-16), 5 surface water COIs (Table 7-40), and 9 TZW COIs (Table 7-43). Risk to birds and mammals from dietary exposure to 19 COIs could not be evaluated (Table 8-30). Risk to amphibians and aquatic plants from 27 COIs (including 19 surface water COIs and 16 TZW COIs) could not be evaluated (Tables 5-11, 5-12, and 6-35 for surface water; Table 6-42 for TZW). As per agreement with EPA (LWG 2010), these COIs are identified as chemicals for which no TRV is available as well as chemicals whose maximum DL exceeded a TRV but whose detected values did not.

Risk assessments are, by design, conservative in the face of uncertainty. However, not all uncertainties create a conservative bias. Some examples of uncertainties that could lead to underestimation of risk include unavailability of exposure or effects data; existing TRVs that might underestimate risk for untested sensitive species; synergistic interactions among the multiple chemicals; and metabolic processes that increase the toxicity of accumulated chemicals.

There is no single 'best' method for summarizing potentially unacceptable ecological risks in a site as large and complex as Portland Harbor, or in a site with as many contaminants posing potentially unacceptable risks. Each reader of the BERA is likely to have need for or interest in different aspects of the risk assessment. Some readers will be interested in chemicals posing potentially unacceptable risks in each medium or sample type; others may be interested in risks to specific assessment or measurement endpoints; while others may be interested in risks within a specific river mile or other geographic locator.

In an effort to make this section as useful to as many readers as possible, the chemicals posing potentially unacceptable risks are sorted and presented in several different ways: by line of evidence, receptor group, assessment endpoint, and by media within each river mile of the Study Area. BERA Attachment 19 provides additional summary details on magnitude and locations of risk not fully presented in this section.

Table 11-1 tallies the COPCs (individual chemicals, sums, or totals) identified as posing potentially unacceptable risk for each assessment endpoint and Table 11-2 provides a more general summary for each ecological receptor group. In total, 89-102 CERCLA contaminants were identified as posing potentially unacceptable risk in this BERA based on $HQ \geq 1$ for at least one receptor-LOE combination. The number of contaminants identified as posing potentially unacceptable risks drops to 74 if all of the individual compound or mixture TRV's for PCB's, DDx (i.e. the various DDD, DDE and DDT compounds) and PAH's are condensed into total PCB, total DDx and total PAH groupings. The maximum HQs and numbers of samples resulting in $HQ \geq 1$ for each

Comment [A2]: Specific table numbers will need to be added here. The individual tables, particularly the table by river mile and medium within each river mile need to be generated. The draft table by assessment endpoint, measurement endpoint and line of evidence has already been shared with LWG, and will be presented as a summary table for Sections 6 through 10, for each major ecological receptor category.

receptor-LOE-COPC combination posing potentially unacceptable risk are presented in Attachment 19:²

- Benthic invertebrates – Eighty-three COPCs were identified via one or more of the sediment, tissue-residue, surface water, and TZW LOEs.³
- Fish – Fifty nine COPCs were identified using the tissue-residue, dietary-dose, surface water, and TZW LOEs.⁴
- Wildlife – Twelve COPCs were identified for birds using the dietary-dose and tissue-residue (egg) LOEs, and six COPCs were identified for mammals using the dietary-dose LOE.
- Amphibians – Thirty-three COPCs were identified using the surface water and TZW LOEs.⁵
- **Aquatic plants** – Thirty-three COPCs were identified using the surface water and TZW LOEs.⁶

Comment [A3]: Will need to retabulate the PUR chemical counts for all of the below bullets.

The spatial extent, magnitude and potential ecological significance of TRV exceedances and the concordance among LOEs were considered to determine risk conclusions for contaminants posing potentially unacceptable risk. The analyses used to draw these conclusions are presented for each receptor group in Sections 6.7, 7.6, 8.3, 9.3, and 10.3. The main conclusions of the BERA by receptor group are briefly summarized below in Section 11.3.

² Counts of COPCs with HQs ≥ 1 are based on HQs derived using [both EPA national ambient water quality criteria \(AWQC\) and the alternative surface water TRVs for total PCBs, 4,4'-DDT, and total DDx derived in this BERA.](#) Both sets of TRVs identify 4,4'-DDT and total DDx as having HQs ≥ 1 in both surface water and TZW. Total PCBs have samples with HQ ≥ 1 in surface water using the AWQC-based TRV, but not the BERA alternative TRV. Neither PCB TRV identifies total PCB as posing potentially unacceptable risk in TZW, as opposed to the AWQC-based TRVs.

³ Eighty-five benthic invertebrate COPCs have HQs ≥ 1 . Petroleum compounds are not CERCLA contaminants, and have been excluded from the final COPC count for sediment and TZW LOEs even though this chemical group may be contributing to potentially unacceptable risk.

⁴ Sixty fish COPCs have HQs ≥ 1 . Petroleum compounds are not CERCLA contaminants and have been excluded from the COPC count for the TZW LOE even though this chemical group may be contributing to potentially unacceptable risk.

⁵ Thirty-four amphibian COPCs have HQs ≥ 1 . Petroleum compounds are not CERCLA contaminants and have been excluded from the COPC count for the TZW LOE even though this chemical group may be contributing to potentially unacceptable risk.

⁶ Thirty-four aquatic plant COPCs have HQs ≥ 1 . Petroleum compounds are not CERCLA contaminants and have been excluded from the COPC count for the TZW LOE even though this chemical group may be contributing to potentially unacceptable risk.

Table 11-1. COPCs with HQ ≥ 1 Organized by Assessment Endpoint and Line of Evidence for the Portland Harbor BERA

Line of Evidence (LOE)	COPCs with HQ ≥ 1
Assessment Endpoint:^a Benthic Invertebrate Survival, Growth, and Reproduction	
Macroinvertebrates (e.g., amphipods, isopods, bivalves, gastropods, oligochaetes, insects, decapods)	
Survival and biomass of <i>Chironomus dilutus</i> and <i>Hyalella azteca</i> exposed to site sediments compared with reference area sediments	Responses based on chemical mixtures; no individual COPCs identified
Concentrations in site sediment compared with effect levels derived from FPM and LRM models (i.e., SQVs) predicting reduced survival or biomass based on Portland Harbor surface sediment concentrations and toxicity reported for both <i>Hyalella</i> and <i>Chironomus</i> endpoints	6 metals, TBT, 19 individual PAHs or group sums, dibutyl phthalate, 3 SVOCs, 2 phenolic compounds, PCBs, 15 individual pesticides or group sums
Concentrations in site sediment compared with national consensus-based SQGs (PECs and related quotients), and effects-based SQGs (PELs, and related quotients)	8 metals, 14 individual PAHs or group sums, 2 PCBs, 9 individual pesticides or group sums
Concentrations in surface water compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx, ^b ethylbenzene, trichlorethene
Concentrations in shallow TZW compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	14 metals, 16 individual PAHs, 3 SVOCs, the pesticides 4,4'-DDT ^b and total DDx, ^b 16 VOCs, gasoline-range hydrocarbons, cyanide and perchlorate
Empirical (field-collected) whole-body concentrations of epibenthic organisms compared with tissue TRVs	None
Steady-state estimates of laboratory-exposed whole-body concentrations in <i>Lumbriculus</i> compared with tissue TRVs	Arsenic, copper, zinc, TBT, PCBs, total DDx
Predicted (BSAF) whole-body concentrations of <i>Lumbriculus</i> compared with tissue TRVs	TBT, PCBs, total DDX
Bivalves (clams, mussels)	
Empirical (field-collected) whole-body concentrations in <i>Corbicula fluminea</i> and freshwater mussels compared with tissue TRVs	Copper, zinc, TBT, PCBs
Steady-state estimates of laboratory-exposed whole-body concentrations in <i>Corbicula fluminea</i> compared with tissue TRVs	TBT, BEHP, total DDx

Table 11-1. COPCs with HQ ≥ 1 Organized by Assessment Endpoint and Line of Evidence for the Portland Harbor BERA

Line of Evidence (LOE)	COPCs with HQ ≥ 1
Predicted (BSAF) whole-body concentrations in <i>Corbicula fluminea</i> compared with tissue TRVs	Total PCBs, total DDx
<i>Corbicula fluminea</i> survival compared with control data from bioaccumulation tests	Responses based on chemical mixtures; no individual COPCs identified
Survival and biomass of <i>Chironomus dilutus</i> and <i>Hyaella azteca</i> exposed to site sediments, compared with reference sediments	Responses based on chemical mixtures; no individual COPCs identified
Concentrations in surface water compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx, ^b ethylbenzene, trichlorethene
Concentrations in shallow TZW compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	14 metals, 16 individual PAHs, 3 SVOCs, the pesticides 4,4'-DDT ^b and total DDx, ^b 16 VOCs, gasoline-range hydrocarbons, ^c cyanide and perchlorate
Concentrations in site sediment compared with national consensus-based SQGs (PECs and related quotients) and effects-based SQGs (PELs and related quotients)	8 metals, 14 individual PAHs or group sums, 2 PCBs, 9 individual pesticides or group sums
Decapods (crayfish)^d	
Empirical whole-body concentrations in crayfish compared with tissue TRVs	Copper
Predicted (BSAF or FWM) whole-body concentrations in crayfish compared with tissue TRVs	Total PCBs, total DDx
Concentrations in site sediment compared with national consensus-based SQGs (PECs and related quotients) and effects-based SQGs (PELs and related quotients)	8 metals, 14 individual PAHs or group sums, 2 PCBs, 9 individual pesticides or group sums
Concentrations in surface water compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx, ^b ethylbenzene, trichlorethene
Concentrations in shallow TZW compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of benthic macroinvertebrate survival, growth, and reproduction	14 metals, 16 individual PAHs, 3 SVOCs, the pesticides 4,4'-DDT ^b and total DDx, ^b 16 VOCs, gasoline-range hydrocarbons, ^c cyanide and perchlorate

Table 11-1. COPCs with HQ ≥ 1 Organized by Assessment Endpoint and Line of Evidence for the Portland Harbor BERA

Line of Evidence (LOE)	COPCs with HQ ≥ 1
Assessment Endpoint:^a Fish Survival, Growth, and Reproduction	
Omnivorous Fish (white sturgeon, largescale sucker^e)	
Empirical whole-body concentrations compared with tissue TRVs	Total PCBs
Dietary dose (including incidental sediment ingestion) compared with dietary TRVs	Copper
Concentrations in surface water compared with state WQS, national AWQC, ^b or effects-based values derived from the literature that are protective of fish survival, growth, and reproduction	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx ^b , 4,4'-DDT ^b , trichloroethene, ethylbenzene, total PCB ^b No COPCs with HQs ≥ 1
Correlation of lesion prevalence with areas of contamination and/or comparison to lesion-based TRVs (if relevant to receptor species) ⁷	Inconclusive for PAHs
Invertivorous Fish (juvenile Chinook salmon,^f peamouth, sculpin)	
Empirical whole-body concentrations compared with tissue TRVs	Copper, lead, total PCBs, total DDx
Predicted (BSAF or FWM) whole-body concentration compared with tissue TRVs (sculpin only)	Total PCBs, total DDx
Dietary dose (including incidental sediment ingestion) compared with dietary TRVs	Cadmium, copper, TBT
Concentrations in surface water compared with state WQS, national AWQC or effects-based TRVs reported in the literature	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx ^b , trichloroethene
Concentrations in shallow TZW compared with state WQS, national AWQC or effects-based TRVs reported in the literature (sculpin only)	14 metals, 16 PAHs, 3 SVOCs, the pesticides 4,4'-DDT ^b and total DDx, ^b 16 VOCs, gasoline-range hydrocarbons, ^c cyanide and perchlorate
Piscivorous Fish (northern pikeminnow, smallmouth bass)	
Empirical whole-body concentrations compared with tissue TRVs	Antimony, lead, total PCBs

Comment [A4]: Footnote will need renumbered to match footnotes in rest of Table 11-1. Renumbering is an editorial change.

⁷ The lesion prevalence line of evidence is not a primary LOE in the BERA, as it does not directly address any BERA assessment endpoint. Since effects on survival, growth or reproduction cannot be quantified from the lesion LOE, no quantitative risk management recommendations can be derived from the lesion LOE.

Table 11-1. COPCs with HQ ≥ 1 Organized by Assessment Endpoint and Line of Evidence for the Portland Harbor BERA

Line of Evidence (LOE)	COPCs with HQ ≥ 1
Predicted (BSAF or FWM) whole-body concentrations compared with tissue TRVs (smallmouth bass only)	This LOE was not evaluated because empirical tissue data were available from all exposure areas.
Concentrations in surface water compared with reported state WQS, national AWQC, ^b or effects-based TRVs reported in the literature	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP
Dietary dose (including incidental sediment ingestion) compared with dietary TRVs	Copper
Detritivorous Fish (Pacific lamprey ammocoete^f)	
Empirical whole-body concentration compared with tissue TRV	Copper
Concentrations in surface water compared with state WQS, national AWQC, or literature-based values that are protective of early life stages.	Zinc, monobutyltin, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx^b, 4,4'-DDT^b, trichloroethene, ethylbenzene, total PCB^b No COPCs with HQs ≥ 1 ^b
Concentration in shallow TZW compared with state WQS, national AWQC, or effects-based values reported in the literature that are protective of early life stages ^g	14 metals, 16 PAHs, 3 SVOCs, the pesticides 4,4'-DDT ^b and total DDx, ^b 16 VOCs, gasoline range hydrocarbons, ^c cyanide and perchlorate
Assessment Endpoint:^a Bird Survival, Growth, and Reproduction	
Invertivorous Birds (spotted sandpiper)	
Dietary dose (including incidental sediment ingestion) compared with dietary TRV	Copper, benzo(a)pyrene, dibutyl phthalate, total PCBs, PCB TEQ, total dioxin/furan TEQ, total TEQ, sum DDE, total DDx, aldrin
Omnivorous Birds (hooded merganser)	
Dietary dose (including incidental sediment ingestion) compared with dietary TRV	Total PCBs
Piscivorous Birds (osprey, bald eagle)	
Dietary-based approach incorporating food chain transfer of contaminants from appropriate fish species (assuming all exposure comes from prey fish) and incidental sediment ingestion	Lead, mercury, total PCBs
Measured concentrations in osprey eggs compared with egg- or embryo-based TRVs for DDT and metabolites, PCBs, and dioxin-like compounds	Total PCBs, PCB TEQ, total dioxin/furan TEQ, total TEQ, sum DDE ^h

Table 11-1. COPCs with HQ ≥ 1 Organized by Assessment Endpoint and Line of Evidence for the Portland Harbor BERA

Line of Evidence (LOE)	COPCs with HQ ≥ 1
Assessment Endpoint:^a Mammal Survival, Growth, and Reproduction	
Aquatic-Dependent Mammals (mink, river otter)	
Dietary dose compared with dietary TRVs	Aluminum, lead, total PCBs, PCB TEQ, total dioxin/furan TEQ, total TEQ
Assessment Endpoint:^a Amphibian Survival, Growth, and Reproduction (frogs, salamanders)	
Concentrations in surface water compared with state WQS, national AWQC, or effects-based values reported in the literature that are protective of sensitive life stages	Zinc, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx ^b
Concentrations in shallow TZW compared with state WQS, national AWQC, or effects-based values reported in the literature that are protective of sensitive life stages	11 metals, 8 PAHs, the SVOC 1,2-dichlorobenzene, the pesticides 4,4'-DDT ^b and total DDx, ^b 8 VOCs, gasoline-range hydrocarbons, ^c and the conventionals cyanide and perchlorate
Assessment Endpoint:^a Aquatic Plant Survival, Growth, and Reproduction (phytoplankton, periphyton, macrophytes)	
Concentrations in surface water compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of sensitive life stages (e.g., germination, emergence, early life stage growth)	Zinc, benzo(a)anthracene, benzo(a)pyrene, naphthalene, BEHP, total DDx ^b
Concentrations in shallow TZW compared with state WQS, national AWQC, or effects-based values derived from the literature that are protective of sensitive life stages (e.g., germination, emergence, early life stage growth)	11 metals, 8 PAHs, the SVOC 1,2-dichlorobenzene, the pesticides 4,4'-DDT ^b and total DDx, ^b 8 VOCs, gasoline-range hydrocarbons, ^c and the conventionals cyanide and perchlorate

^a The assessment endpoints for all receptors are based on protection and maintenance of their populations and the communities in which they live, except that the health of threatened or endangered species is to be protected at the level of the individual organism. Per the SOW, EPA's Problem Formulation (Attachment 2), and as stated in the Programmatic Work Plan (Integral et al. 2004b), the assessment endpoints were expressed as the survival, growth, and reproduction of each receptor group.

^b Risk estimates for total PCBs, 4,4'-DDT, and total DDx for the surface water and TZW LOEs are based [HQS derived using both EPA national ambient water quality criteria \(AWQC\) and the alternative surface water TRVs for total PCBs, 4,4'-DDT, and total DDx derived in this BERA. Both sets of TRVs identify 4,4'-DDT and total DDx as having HQs ≥ 1 in both surface water and TZW. Total PCBs and 4,4'-DDT have samples with HQ ≥ 1 in surface water using the AWQC-based TRV, but not the BERA alternative TRV. Neither PCB TRV identifies total PCB as posing potentially unacceptable risk in TZW, on the alternative total PCBs and 4,4'-DDT TRVs for protection of directly exposed aquatic organisms, rather than the selected AWQC-based TRVs. Additional exceedances occur using the AWQC-based TRVs and HQs, as presented in the surface water and TZW risk characterization sections for each receptor group.](#) The alternative TRVs are considered more appropriate for evaluating direct exposure of aquatic organisms because the [national](#) AWQC are based on protection of dietary risks to mammals and birds.

- ^c ~~The HQ for gasoline-range hydrocarbons is ≥ 1 ; however the COPC was not included in the counts of COPCs with HQs ≥ 1 because counts are based only on CERCLA contaminants.~~
- ^d Although these LOEs are components of the benthic invertebrate community, the bivalve population and decapod population assessment endpoints are presented separately in this table. Evaluation of sediment toxicity to *Chironomus* and *Hyalella* and comparison of surface water and shallow TZW concentrations to TRVs were each conducted and presented only once as part of the benthic invertebrate community assessment. Similarly, comparison of sediment concentrations to published SQGs also occurred and was presented only once as part of the benthic community assessment.
- ^e Carp is not a receptor of concern for the BERA but whole-body carp tissue was analyzed for dioxin-like chemicals, including PCB congeners; for these chemicals, carp is a surrogate for other omnivorous fish species.
- ^f Juvenile Chinook salmon and Pacific lamprey ammocoetes were evaluated at the organism level because they have special status are (juvenile Chinook is federally threatened and Pacific lamprey is an Oregon state sensitive species of special concern to Tribes); effect thresholds based on reproduction are used as a surrogate for growth in juvenile Chinook salmon and Pacific lamprey ammocoetes.
- ^g The TZW exposure pathway for fish receptors is considered complete and significant for only sculpin and lamprey ammocoetes. The ecological CSM shows a complete TZW exposure pathway for sucker, carp, and sturgeon but categorizes the pathway as insignificant.
- ^h Bald eagle only based on extrapolation from osprey eggs and comparison to a NOAEL-based TRV. For osprey, all HQ < 1.
- | | | |
|--|--|---|
| AWQC – ambient water quality criteria | EPA – US Environmental Protection Agency | SVOC – semivolatile organic compound |
| BEHP – bis(2-ethylhexyl) phthalate | FWM – food web model | TBT – tributyltin |
| BERA – baseline ecological risk assessment | HQ – hazard quotient | TEQ – toxic equivalent |
| BSAF – biota-sediment accumulation factor | LOE – line of evidence | total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT) |
| CERCLA – Comprehensive Environmental Response, Compensation, and Liability Act | PAH – polycyclic aromatic hydrocarbon | TRV – toxicity reference value |
| COPC – contaminant of potential concern | PCB – polychlorinated biphenyl | TZW – transition zone water |
| CSM – conceptual site model | PEC – probable effects concentration | VOC – volatile organic compound |
| DDD – dichlorodiphenyldichloroethane | PEL – probable effects level | WQS – water quality standards |
| DDE – dichlorodiphenyldichloroethylene | SOW – scope of work | |
| DDT – dichlorodiphenyltrichloroethane | SQG – sediment quality guideline | |
| | SQV – sediment quality value | |

Comment [A5]: Footnote C need deleted, and the remaining footnotes relettered. EPA considers the TPH fractions to be CERCLA contaminants.

Table 11-2. Contaminants Posing Potentially Unacceptable Risk Organized by Receptor Group

COPC ^a	Benthic Invertebrates	Fish	Birds	Mammals	Amphibians	Aquatic Plants
Metals						
Aluminum				X		
Antimony		X				
Arsenic	X					
Barium	X	X			X	X
Beryllium	X	X				
Cadmium	X	X			X	X
Cobalt	X	X				
Copper	X	X	X		X	X
Iron	X	X			X	X
Lead	X	X	X	X	X	X
Magnesium	X	X			X	X
Manganese	X	X			X	X
Mercury		X	X			
Nickel	X	X			X	X
Potassium	X	X			X	X
Sodium	X	X			X	X
Vanadium	X	X				
Zinc	X	X			X	X
Butyltins						
Monobutyltin	X	X				
Tributyltin	X	X				
PAHs						
2-Methylnaphthalene	X	X			X	X
Acenaphthene	X	X			X	X
Anthracene	X	X			X	X
Benzo(a)anthracene	X	X			X	X
Benzo(a)pyrene	X	X	X		X	X
Benzo(b)fluoranthene	X	X				
Benzo(g,h,i)perylene	X	X				
Benzo(k)fluoranthene	X	X				
Chrysene	X	X				
Dibenzo(a,h)anthracene	X	X				
Fluoranthene	X	X				

Table 11-2. Contaminants Posing Potentially Unacceptable Risk Organized by Receptor Group

COPC ^a	Benthic Invertebrates	Fish	Birds	Mammals	Amphibians	Aquatic Plants
Fluorene	X	X			X	X
Ideno(1,2,3-cd) pyrene	X	X				
Naphthalene	X	X			X	X
Phenanthrene	X	X			X	X
Pyrene	X	X				
Phthalates						
BEHP	X	X			X	X
Dibutyl phthalate			X			
SVOCS						
1,2-Dichlorobenzene	X	X			X	X
1,4-Dichlorobenzene	X	X				
Dibenzofuran	X	X				
PCBs						
Total PCBs	X	X	X ^c	X		
PCB TEQ			X ^c	X		
Dioxins/furan TEQ			X ^c	X		
Total TEQ			X ^c	X		
VOCs						
1,1-Dichloroethene	X	X				
1,2,4-Trimethylbenzene	X	X			X	X
1,3,5-Trimethylbenzene	X	X				
Benzene	X	X				
Carbon disulfide	X	X			X	X
Chlorobenzene	X	X			X	X
Chloroethane	X	X			X	X
Chloroform	X	X			X	X
cis-1,2-Dichloroethene	X	X				
Ethylbenzene	X	X			X	X
Isopropylbenzene	X	X			X	X
Toluene	X	X			X	X
Trichloroethene	X	X				
m,p-Xylene	X	X				
o-Xylene	X	X				
Total xylenes	X	X				

Table 11-2. Contaminants Posing Potentially Unacceptable Risk Organized by Receptor Group

COPC ^a	Benthic Invertebrates	Fish	Birds	Mammals	Amphibians	Aquatic Plants
Pesticides						
Aldrin			X			
4,4'-DDD	X					
sum DDE			X			
4,4'-DDT	X	X			X	X
Total DDx	X	X	X		X	X
Other Chemicals						
Cyanide	X	X			X	X
Perchlorate	X	X			X	X

^a The COPCs listed in this table are CERCLA contaminants. Several additional contaminants may also contribute to potentially unacceptable risk. These contaminants include TPH, ammonia, and sulfides.

BEHP – bis(2-ethylhexyl) phthalate

CERCLA – Comprehensive Environmental Response,
Compensation, and Liability Act

COPC – chemical of potential concern

DDD – dichlorodiphenyldichloroethane

DDE – dichlorodiphenyldichloroethylene

DDT – dichlorodiphenyltrichloroethane

HCH – hexachlorocyclohexane

HQ – hazard quotient

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

TEQ – toxic equivalent

total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD,
2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT)

Comment [A6]: TPH is a CERCLA contaminant, and the appropriate fractions should be listed as such under surface water and sediment in Table 11-2.

11.2 BACKGROUND AND UPRIVER CONCENTRATIONS

For all contaminants posing potentially unacceptable risk, Attachment 11 presents a comparison of background and Study Area 95th percentile UCLs in sediment and surface water. For aluminum, dibutyl phthalate, benzyl alcohol, and alpha-endosulfan, background sediment UCLs are the same as or higher than Study Area UCLs. The background surface water UCL concentration is higher than the Study Area UCL only for aluminum. Attachment 11 also includes a comparison of concentrations in fish tissue from the upriver reach and the Study Area for all fish tissue-residue and wildlife dietary contaminants posing potentially unacceptable risk. Although fish tissue data from the upriver reach are insufficient to allow calculation of UCLs, their concentrations are similar to those in the Study Area for aluminum, mercury, and copper, as presented in Section 7.1.5.

Background concentrations for sediment and surface water, and upriver concentrations for fish tissue provide context for Study Area risk predictions but were not used to discount risks or influence risk estimates. Where background concentrations exceed screening-level TRVs or upriver fish tissue concentrations exceed tissue TRVs, upriver or regional sources may be contributing to unacceptable risks in the Study Area.

11.3 ECOLOGICAL RISK CONCLUSIONS

The risk conclusions across all ecological receptor groups are combined and briefly summarized in this section to provide a general overview of ecological risks and to identify the receptor-COPC pairs that, given the magnitude and extent of risk, are reasonably likely to result in adverse effects on the assessment endpoints that were selected in the Problem Formulation to represent the valued ecological attributes of the Study Area. In other words, what is the ecological significance of the risks identified in Sections 6 through 10? To reiterate, the analyses used to draw these conclusions are presented in Sections 6.7, 7.6, 8.3, 9.3, and 10.3, and are not repeated here. For example, this section (11.3) contains statements with qualitative adjectives like “limited” or “moderate” when describing the spatial extent of exposure to a COPC at concentrations yielding HQs ≥ 1 . Statements such as, “uncertainty in the tissue-residue TRV is more likely to over- than underpredict risk” are made without repeating the supporting evidence. EPA (1997) ecological risk assessment guidance calls for professional judgement on the part of risk assessors when evaluating the ecological significance of identified risks. In cases such as these, the reader interested in the details should refer back to the risk conclusions section for the relevant receptor group. The main conclusions of the BERA by receptor group are presented in Sections 11.3.1 through 11.3.5. Section 11.3 closes with a brief synopsis of potential future benthic community risks in erosional sections of the Study Area.

Comment [A7]: A discussion of ecological significance of risks needs to be appended as a new subsection to the end of Section 3.3 (assessment endpoint selection). The criteria for ecological significance EPA described in the draft executive summary can be expanded on in this new section. If the text added to Section 3.3 does a good job defining ecological significance, the text in Section 11.3 should largely be acceptable as written, with the exception of the last paragraph of Section 11.3.3.

11.3.1 Benthic Invertebrate Community

COPCs occur at concentrations that are projected to pose unacceptable benthic risks for about 7% of the Study Area. Based on GIS estimates extrapolated from the empirical sediment toxicity data, Level 2 (moderate) and Level 3 (severe) toxicity combined are projected to extend over between 4 – 8% of the area of surface sediment area within the Study Area. Unlike other ecological receptors, for which risk was evaluated on a chemical-specific basis, risk to the benthic invertebrate community was evaluated in large part by considering exposure to the mixture of chemicals present in the Study Area sediments, using toxicity tests and multivariate predictive models based on the toxicity test results. Point-by-point assessment of potential effects on benthic organisms using data from toxicity testing, modeling, and benthic tissue-residue analyses indicates that metals, TBT, PAHs, several SVOCs, two phenolic compounds, dibutyl phthalate, total PCBs, total DDx, and other pesticides pose potentially unacceptable risk. Several other contaminants (TPH, ammonia, and sulfides) may also contribute to potentially unacceptable risk at some areas. A WOE was assessed to identify contaminants that were most likely posing unacceptable risk. Based on that evaluation, the primary COPCs in sediment that likely pose potentially unacceptable risk to the benthic community or populations are PAHs, PCBs, and total DDx. Although other contaminants may also contribute to unacceptable risk, their distribution and magnitude of risk tends to be represented by the distribution and magnitude of primary COPCs. One exception is the certain contaminants associated with the localized TZW investigation areas. In these areas, VOCs, cyanide, and perchlorate may also pose potentially unacceptable risks; however, these contaminants often co-occur with PAHs and DDx.

The phenolic compound 4-methylphenol may also be contributing to benthic community risk. The analysis conducted for the BERA shows that the sediment exposure pathway is sufficient to be of concern for 4-methylphenol. Widely distributed throughout the Study Area, this contaminant is found in both contaminated and otherwise uncontaminated areas. Methylated phenols are readily biodegraded under aerobic conditions, and 4-methylphenol is expected to have a half-life in sediment on the order of days. That 4-methylphenol was found suggests the presence of ongoing sources; however, whether and to what extent the source is degradation of historical contamination versus influx from ongoing point or non-point discharges is not known.

Sediment profile images of the surface sediment suggest that the physical environment (sediment grain size, transport regime, bottom slope) in the Study Area can explain the presence of early colonizing, transitional, and mature benthic communities in 90% of the images evaluated. In these cases, the successional stage matched the expected community structure based on the physical regime and habitat characteristics. In the vast majority of cases, mature benthic communities occurred in fine-grained depositional environments; early colonizing or transitional communities were found in less physically stable areas (for example, with steep slopes, active sediment transport, high rates of deposition, or physical disturbance). In the 31 (of 377) cases where the community successional stage was not as might be predicted by the physical environment, about two-thirds (19) occur between RM 5.0 and RM 9.0. The greatest combined area associated with potentially unacceptable risk to the benthic community was found in this same reach, suggesting possible chemical toxicity, among other potential factors, as the reason for the presence of lower successional stages. These qualitative results suggest that overall, the benthic community in the Study Area is typical of a large river system that is strongly influenced by physical processes. Impacts from sediment contamination appear to be limited to certain depositional areas that have received historical releases of contamination.

11.3.2 Fish

The fish assessment endpoints are survival, growth and reproduction of omnivorous, invertivorous, and piscivorous fish, as well as survival and growth of detritivorous fish. The assessment endpoints are based on protection and maintenance of populations and the communities in which they live, except for Pacific lamprey ammocoete and juvenile Chinook salmon, which, as special status species, are to be protected at the organism level.

Total PCBs were found to pose low risk to populations of piscivorous fish and the small-home-range invertivorous fish sculpin. Total PCB tissue-residue HQs ≥ 1 were calculated for smallmouth bass, northern pikeminnow, and sculpin samples from locations throughout the Study Area (max HQ = 9.4). HQs < 1 for juvenile Chinook salmon and peamouth show that risk to sculpin does not imply risk to invertivorous fish with larger home ranges. Together, the low Study Area-wide tissue-residue HQ of 1.6 for largescale sucker in combination with HQs < 1 for most omnivorous fish samples and with uncertainty in effects data indicate that risk to omnivorous fish is negligible.

The potential for adverse effects on all of the fish assessment endpoints from total PCBs was assessed to be low: the other LOE for PCBs—surface water—resulted in HQs < 1,⁸ tissue-residue HQs ≥ 1 occurred over only a moderate spatial extent (or in relatively few samples for large-home-range fish), and uncertainty in the tissue-residue TRV is more likely to overpredict than underpredict risk. Uncertainties in the fish risk assessment are described in more detail in Sections 7.1.4, 7.2.5.1 and 7.2.5.4. The tissue-residue TRV for total PCBs is conservative because it is based partially on uncertain toxicity data, including field data from contaminated sites where other contaminants were also present, suggesting that the TRV reflects toxicity from chemicals other than PCBs.

The spatial extent of dietary risk to juvenile Chinook salmon from cadmium encompasses a substantial portion of the Study Area. However, the assumption that juvenile Chinook consume benthic invertebrates, rather than the pelagic prey they are known to eat, overestimates exposure. The selected TRV also very likely overestimates risk because it is 3 orders of magnitude below the lowest salmon-specific NOAEL.

The spatial extent of dietary and tissue-residue risk from copper to several fish (sculpin, juvenile Chinook salmon, Pacific lamprey ammocoetes, northern pikeminnow, largescale sucker, and juvenile white sturgeon) also encompasses a substantial portion of the Study Area. The copper-fish TRVs are highly uncertain. The dietary TRV could not be replicated in subsequent studies, and the tissue-residue TRV is within the range of copper nutritional requirements for some (but not all) fish species. Furthermore, predictions of risk to fish based on tissue concentrations copper is highly uncertain because fish regulate this essential metal.

Several COPCs in TZW were identified as posing risk to individual fish, but not their populations. Benthic fish, including burrowing fish (lamprey ammocoetes) and fish that feed on benthic organisms (sculpin), have relatively low exposure to porewater compared with surface water because of their feeding habits and respiratory requirements. Risks from elevated TZW contaminant concentrations are localized to the portion of the local population exposed to the discharge area of the contaminated groundwater plumes. Portions of the local population not exposed to TZW in the groundwater discharge area are likely unaffected by the TZW contaminants. Although as TZW samples were collected from only a limited number of locations in the Study Area, the total proportion of the site with groundwater discharge into the Willamette River, as well as contaminant concentrations in TZW from any such additional areas, are unknown. For this reason concentrations of COPCs in shallow TZW likely overestimate exposure, to an uncertain degree. Because TZW exceedances are localized, none-the potential of the TZW COPCs is likely to pose risk to Study Area benthic invertebrate or fish populations is also uncertain. However, 38 TZW COPCs,⁹ 6 metals (barium, iron, manganese, sodium,

Comment [A8]: Recheck this count.

⁸ When calculated using the alternative TRV for protection of directly exposed aquatic organisms rather than the AWQC, which is based on protection of mink through dietary exposure.

⁹ Petroleum hydrocarbons were evaluated as an uncertainty and gasoline-range aliphatic hydrocarbons (C10-C12) have HQ > 10 over a limited spatial extent and also pose potentially unacceptable risk to individual lamprey.

vanadium, and zinc), 16 PAHs (2-methylnaphthalene, acenaphthene, anthracene, benzo(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, chrysene, dibenzo(a,h)anthracene, fluoranthene, fluorene, indeno(1,2,3-cd)pyrene, naphthalene, phenanthrene, pyrene), 2 SVOCs (1,2-dichlorobenzene, 1,4-dichlorobenzene), the pesticides 4,4'-DDT and total DDX, 10 VOCs (benzene, carbon disulfide, chlorobenzene, chloroform, cis-1,2-dichloroethene, ethylbenzene, o-xylene, toluene, total xylenes and trichloroethene), cyanide and perchlorate have high concentrations in localized areas that could adversely affect Pacific lamprey ammocoetes at those locations. The magnitude of risk to individual lamprey from these COPCs is unknown however, because the TRVs were derived to be protective of the most sensitive species and are likely to overpredict risk to Pacific lamprey which has been shown to have average or lower sensitivity than most aquatic species for several chemicals causing toxicity from different modes of action (Andersen et al. 2010). Three of the 38 COPCs (excluding petroleum hydrocarbons.) with HQs > 10 are naturally occurring metals (barium, iron, and manganese) and there is substantial uncertainty as to whether their source is anthropogenic.

Risk to fish from other COPCs that resulted in HQs ≥ 1 in the final step of the risk characterization were found unlikely to result in ecologically significant adverse effects on the fish populations. The rationale for this conclusion is that TZW exposure assumptions likely overestimate risk, the TRV overestimates risk, and the great majority of samples result in HQs < 1 (indicating a limited spatial extent of potentially unacceptable risk).

11.3.3 Wildlife

The avian assessment endpoints are survival, growth, and reproduction of invertivorous, omnivorous, and piscivorous birds. The mammalian assessment endpoint was survival, growth, and reproduction of aquatic-dependent mammals. The assessment endpoints are based on protection and maintenance of populations and the communities in which they live, except for threatened or endangered species, which are to be protected at the organism level.

Total PCBs pose the primary risk. Mink and river otter HQs ≥ 1 throughout the Study Area (mink HQ = 19 to 33, river otter HQ = 21 to 31) indicate that PCBs pose ecologically significant risk of reduced reproductive success to populations of both receptors in the Study Area. While this BERA has established that PCBs pose the potential for adverse effects, the true effect of PCB exposure on Study Area populations is still unknown because of a number of uncertainties. These include quantifiable uncertainties about dietary exposure and about PCB dose-response, and quantifiable uncertainty about the level of effect associated with a population-level response. These uncertainties have not been fully examined in the BERA. [The known sensitivity of mink to PCBs is demonstrated in EPA's AWQC for PCBs, which is based on protection of dietary toxicity of bioaccumulated PCBs in prey of the aquatic-dependent mink, not on PCB toxicity to fish, aquatic invertebrates or plants. This sensitivity of mink to PCBs is likely](#)

the reason the BERA alternative water column TRV for total PCB is higher than the EPA AWQC.

Reproductive success in spotted sandpiper, bald eagle, and osprey might also be reduced because of PCB exposure, as indicated by spotted sandpiper and bald eagle HQs ≥ 1 throughout the Study Area (max HQ = 12 for sandpiper and 3.9 for eagle) and by less widespread osprey HQs ≥ 1 (max HQ = 4.4). Overall, a greater degree of uncertainty is associated with PCB risk estimates for birds than for mammals because of uncertainty about exposure and uncertainty in the effects data. Uncertainty is higher for otter than for mink because otter-specific effects data are lacking.

Comment [A9]: Bald eagle risks will have to be recalculated using LOAEL TRVs in Section 8, because of the change in status of bald eagle from threatened to not threatened. The new results will have to be summarized here. For dietary dose of PCB, the LOAEL TRV is 2x the NOAEL TRV, meaning, for example, the HQ = 3.9 given here will reduce to HQ = 1.9 or 2.0 depending on rounding.

Total TEQ exposure also poses ecologically significant risk of reduced reproductive success to populations of mink (with HQs up to 12). Total TEQ risk to birds and otter is low, considering the WOE for eagle and osprey, the more limited spatial extent of TRV exceedances for sandpiper, and the low magnitude of HQs for river otter. PCBs are responsible for the majority of total TEQ exposure, in that PCB TEQ HQs generally constitute the majority of the total TEQ HQs. For example, mink total TEQ HQs are ≥ 1 in 16 of 109 potential prey samples; of these samples, PCB TEQ HQs are ≥ 1 in 15 samples and total dioxin/furan TEQ HQs are ≥ 1 in only 4 samples. As is the case for total PCBs, a greater degree of uncertainty is associated with total TEQ risk estimates for birds and otter than for mink because of uncertainties in both exposure and effects data for birds and uncertainty in effects data for otter.

Osprey egg data indicate that DDX compounds pose negligible risk to osprey and low to negligible risk of reduced reproductive success to individual bald eagles within limited portions of the Study Area. The only other wildlife receptor with a DDX HQ ≥ 1 is the spotted sandpiper. DDX compounds pose negligible risk to the spotted sandpiper population because the HQs are of low magnitude, span a limited spatial extent, and based on uncertainties in exposure and effects that likely cause overestimates of risk.

The spatial extent of copper HQ ≥ 1 in sandpiper encompasses a large portion of the Study Area; however, risk is negligible. Only one prey item (laboratory-exposed worms) had tissue concentrations associated with an HQ ≥ 1 . Copper HQs based on a mixed-species diet are < 1 . Additionally, the selected TRV was below the lowest bounded literature-reported NOAEL for birds.

Risk to wildlife from other COPCs that resulted in HQ ≥ 1 in the final step of the risk characterization were found unlikely to result in ecologically significant adverse effects on the receptor populations because the HQs are of low magnitude, span a limited spatial extent, and are based on uncertainties in exposure and effects that likely cause an overestimate of risk. The combined toxicity of dioxins/furans and dioxin-like PCBs, expressed as total toxic equivalents (TEQ), poses the potential risk of reduced reproductive success in mink, river otter, spotted sandpiper, bald eagle and osprey. The PCB TEQ fraction of the total TEQ is responsible for the majority of total TEQ exposure, but the total dioxin/furan TEQ fraction also exceeds its TRV in some locations of the Study Area.

Comment [A10]: Text slightly edited from LWG's executive summary, retained in EPA exec. Summary rewrite.

11.3.4 Amphibians

The amphibian assessment endpoints are survival, growth, and reproduction of amphibians. The assessment endpoints are based on protection and maintenance of populations and the communities in which they live, except for threatened or endangered species, which are to be protected at the organism level. For all COPCs with HQs ≥ 1 , the risk to amphibian populations was assessed to be negligible. COPCs in surface water samples resulting in HQ ≥ 1 were found at concentrations below amphibian-specific thresholds or were collected during non-reproductive periods (when amphibians may not be present in the Study Area). For the TZW LOE, the great majority of samples result in HQs < 1 , indicating limited spatial extent of exceedance. Although risk to amphibians from TZW is highly uncertain, it is likely to be negligible because significant exposure to Study Area TZW by this receptor group is unlikely.

11.3.5 Aquatic Plants

The aquatic plant assessment endpoints are survival, growth, and reproduction of aquatic plants. The assessment endpoints are based on protection and maintenance of populations and the communities in which they live, except for threatened or endangered species, which are to be protected at the organism level. For all COPCs with HQs ≥ 1 , the risk to aquatic plant populations was assessed to be negligible. The same COPCs whose surface water HQ is ≥ 1 were found in the great majority of samples to have HQ < 1 and at concentrations generally below algae-specific thresholds. For the TZW LOE, the great majority of samples result in HQs < 1 , indicating limited spatial extent of exceedance.

11.3.6 Potential Future Risks to the Benthic Community

Risk to the benthic community was assessed both for current conditions in the Study Area and estimated future conditions. The future condition assessment is based on the maximum bed change scenario presented in the draft RI (Map 3.4-7) and a sample-by-sample evaluation of changes in status of predicted risk in the erosional areas based on comparison to site-specific SQVs. Attachment 18 presents the approach and results of the current and future risk predictions in the erosional areas of the Study Area. For the majority of erosional sediments (approximately 60%), there was no change of status in predicted risk to the benthic community (i.e., the sediment quality was similar at the erosional depth and the surface). This finding is not surprising because the erosional sediments are predicted to be primarily sands. Of the remaining erosional sediments, approximately 24% is predicted to be more contaminated in the future. The last 16% of the erosional area is predicted to be cleaner after the erosional event.

12.0 RISK MANAGEMENT RECOMMENDATIONS

This section presents the LWG's ecological risk management recommendations to develop and evaluate remedial alternatives that are protective of ecological resources. Risk management recommendations are provided in four main parts:

- Section 12.1 presents recommended COCs for populations of fish and wildlife receptors.¹⁰
- Section 12.2 presents recommendations regarding contaminants present in TZW. TZW risk management recommendations are presented separately from those for other exposure media because the TZW LOE focuses on a spatially limited set of nine TZW sampling areas; the other exposure media (sediment, tissues, and surface water) were evaluated Study Area-wide. Furthermore, the TZW sampling areas were selected to capture information at locations with known or likely pathways for ongoing sources (discharge of upland contaminated groundwater), whereas the other exposure media were investigated because they represent complete exposure pathways to ecological receptors from contaminated sediment. Thus, both the nature and extent of risk as well as the alternatives for addressing them are unique for TZW.
- Section 12.3 presents risk management recommendations for protection of the benthic invertebrate community. As the BERA's benthic risk conclusions rely heavily on LOEs that do not identify specific COPCs (i.e., empirical measurements of sediment toxicity, predictions of sediment toxicity based on multivariate statistical models, and benthic community data from SPI imagery), this section recommends methodologies for delineating benthic AOCs and for evaluating the degree to which remedial action alternatives protect the benthic community.
- Section 12.4 summarizes the risk management recommendations.

12.1 Recommendation of COCs for Study Area Populations of Fish and Wildlife Receptors

In this section, the entire set of contaminants identified as posing potentially unacceptable risk to fish and wildlife receptors is evaluated. The purpose of the evaluation is to identify the COPCs for fish and wildlife receptors to use in the FS to develop and evaluate remedial alternatives that are protective of ecological resources. This subset of COPCs constitutes the recommended COCs.

As discussed in Section 3, the assessment endpoints for most of the ecological receptors identified in EPA's Problem Formulation are for protection of the populations of fish, birds, mammals, and amphibians, and for protection of communities of benthic invertebrates and aquatic plants. The exceptions are that assessment endpoints for special

¹⁰ Where secondary benthic LOEs support these recommendations for fish and wildlife receptors, they are identified.

status species identified in EPA's Problem Formulation (i.e., bald eagle, juvenile Chinook salmon, and Pacific lamprey ammocoetes) are for protection at the level of the organism.

The COC recommendations provided in this section are intended to address Study Area-wide risks to receptor populations. These recommendations are also intended to be protective of the aquatic plant community and receptors assessed at the organism level, except risk to Pacific lamprey ammocoetes from TZW exposure. Recommendations regarding risks from exposure to contaminants posing potentially unacceptable risk in TZW are presented in Section 12.2. Recommendations regarding identification of benthic risk areas and related protectiveness are provided in Section 12.3.

The remainder of Section 12.1 is presented in three main parts:

- Section 12.1.1 presents the rationale for COC recommendations.
- Section 12.1.2 applies that rationale to recommend COCs.
- Section 12.1.3 provides additional recommendations for the contaminants posing potentially unacceptable risk that are recommended as COCs. This includes recommendations about which receptors of concern should be considered along with the COCs to assess the protectiveness of potential remedies in the FS analysis of alternatives.

12.1.1 Rationale for COC Recommendations

COCs will be used to develop and evaluate remedial alternatives that are protective of ecological resources. The FS will also evaluate whether remedial alternatives for these COCs address the full list of contaminants posing potentially unacceptable risk.

The COC recommendations took into account one or more of the following factors:

- How often, where, and in which media risk thresholds were exceeded
- The ecological relevance (strengths and weaknesses) of the exposure estimates used to calculate HQs
- The toxicological effects associated with the TRV
- The magnitude of the exceedance
- Whether a relationship was found between COPC concentrations in co-located sediment and tissue concentrations (for small-home-range species)
- The relative strength and concordance among LOEs used to evaluate risks
- Comparison of Study Area concentrations with available background or upriver data

Some of these factors are strongly risk-based (e.g., the toxicological effects associated with the TRV, and the relative strength and concordance among LOEs), whereas others

are more directly related to practical FS considerations (e.g., whether a relationship was found between COPC concentrations in co-located sediment and tissue concentrations for small-home-range species, and comparison with available background or upriver data).

12.1.2 COC Recommendations

Table 12-1 summarizes the contaminants posing potentially unacceptable risk in this BERA and whether they are recommended as COCs for fish and wildlife receptors. Contaminants posing potentially unacceptable risk based on the TZW LOE are discussed in Section 12.2. Areas and contaminants posing potentially unacceptable risk to the benthic community are discussed in Section 12.3; however, where benthic tissue-residue and surface water LOEs support the selection of COCs for protection of fish and wildlife, they are noted. Nineteen COPCs with at least one $HQ \geq 1$ have been identified in this BERA for fish and wildlife receptors.^{11,12} The set consists of seven metals, two butyltins, three PAHs, two phthalates, PCBs, dioxins/furans, two pesticides, and one VOC. The specific rationale for COC recommendations—based on the seven factors identified in Section 12.1—follows Table 12-1.

Table 12-1. COC Recommendations for All Receptor Group-LOE Pairs with an $HQ \geq 1$

COPC	Receptor Group-LOE Pairs Resulting in $HQ \geq 1$
Contaminants Recommended as COCs	
PCBs	
Total PCBs	Benthic invertebrate – tissue residue (clam, worm) Fish – tissue-residue (sucker, sculpin, bass, pikeminnow) Mammal – diet (mink, river otter) Bird – diet (sandpiper, osprey, bald eagle, merganser) Bird – tissue-residue (osprey, bald eagle)
Dioxins/Furans	
Total TEQ ^a	Mammal – diet (mink, river otter) Bird – diet (sandpiper) Bird – tissue residue (osprey, bald eagle)
Contaminants Not Recommended as COCs	
Inorganic Metals	
Aluminum	Mammal – diet (mink)

¹¹ PCB TEQ and dioxin/furan TEQ are not included in this count because they are components of the total TEQ.

¹² Risk management recommendations for the benthic community assessment endpoints and the TZW LOE are handled separately and are not included in this COPC count.

Table 12-1. COC Recommendations for All Receptor Group-LOE Pairs with an HQ \geq 1

COPC	Receptor Group-LOE Pairs Resulting in HQ \geq 1
Antimony	Fish – tissue residue (bass)
Arsenic	Benthic invertebrate – tissue residue (worm)
Cadmium	Fish – diet (sculpin, Chinook)
Copper	Benthic invertebrate – tissue residue (clam, crayfish, worm) Fish – diet (sucker, sturgeon, Chinook, peamouth, sculpin, pikeminnow) Fish – tissue-residue (sculpin, Chinook, lamprey, pikeminnow)
	Birds – diet (sandpiper)
Lead	Fish – tissue-residue (peamouth, bass) Birds – diet (osprey) Mammals – diet (mink)
Zinc	Benthic invertebrates – surface water, benthic invertebrate tissue residue (clam, mussel, worm) Fish – surface water (sculpin, bass, pikeminnow) Amphibians – surface water Aquatic plants – surface water
Organometals	
Mercury	Fish – diet (sculpin)
Monobutyltin	Benthic invertebrates – surface water Fish – surface water (sculpin, bass, pikeminnow) Birds – diet (bald eagle)
TBT	Benthic invertebrate (clam and worm tissue residue) Fish – diet (sculpin)
PAHs	
Benzo(a)anthracene	Benthic invertebrates – surface water Fish – surface water (sculpin, bass, pikeminnow) Amphibians – surface water Aquatic plants – surface water
Benzo(a)pyrene	Benthic invertebrates – surface water Fish – surface water (sculpin, bass, pikeminnow) Birds – diet (sandpiper) Amphibians – surface water r Aquatic plants – surface water

Table 12-1. COC Recommendations for All Receptor Group-LOE Pairs with an HQ \geq 1

COPC	Receptor Group-LOE Pairs Resulting in HQ \geq 1
Naphthalene	Benthic invertebrates – surface water Fish – surface water (sculpin, bass, pikeminnow) Amphibians – surface water Aquatic plants – surface water
Phthalates	
BEHP	Benthic invertebrates – surface water, tissue residue (worms) Fish – tissue residue (sculpin, bass,) Fish – surface water (sculpin, bass, pikeminnow) Amphibians – surface water Aquatic plants – surface water
Dibutyl phthalate	Birds – diet (sandpiper)
Pesticides	
Aldrin	Birds – diet (sandpiper)
Total DDx ^b	Benthic invertebrates – surface water, tissue residue (clam, worm) Fish – tissue residue (sculpin) Fish – surface water (sculpin) Birds – diet (sandpiper) Birds – tissue residue (bald eagle) Amphibians – surface water Aquatic plants – surface water
4,4'-DDD	Benthic invertebrate – tissue residue (worms)
VOCs	
Ethylbenzene	Benthic invertebrates – surface water
Trichloroethene	Benthic invertebrates – surface water Fish – surface water (sculpin)

^a Total TEQ includes risk estimates for PCB TEQ and total dioxin/furan TEQ.

^b Total DDx includes risk estimates for the additional DDx components that were also evaluated independently (sum DDE, 4,4'-DDE, and 4,4'-DDT). Risk estimates for the surface water LOE are based on the alternative 4,4'-DDT TRVs for protection of directly exposed aquatic organisms, rather than the AWQC-based TRV. The alternative TRV is considered more appropriate for evaluating direct exposure of aquatic organisms because the AWQC is based on protection of dietary risks to birds.

AWQC – ambient water quality criterion

BEHP – bis(2-ethylhexyl) phthalate

COC – contaminant of concern

COPC – contaminant of potential concern

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

TBT – tributyltin

DDD – dichlorodiphenyldichloroethane
DDE – dichlorodiphenyldichloroethylene
DDT – dichlorodiphenyltrichloroethane
HQ – hazard quotient
LOE – line of evidence

TEQ – toxic equivalent
total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT)
TRV – toxicity reference value
VOC – volatile organic compound

12.1.2.1 Recommended COCs

PCBs

Total PCBs is recommended as a COC because exposure poses a risk of ecologically significant adverse effects to mink and river otter populations. It also poses risk of ecologically significant adverse effects to spotted sandpiper, osprey, sculpin, and smallmouth bass populations and risk of adverse effects to bald eagles. The benthic tissue-residue LOE also supports the selection of PCBs as a COC. These additional risks are lower than the risk to mink and river otter populations. Further risk management recommendations regarding total PCBs are provided in Section 12.1.3.

Dioxins/Furans

Total TEQ is recommended as a COC because exposure poses a risk of ecologically significant adverse effects to mink populations. Total TEQ also poses risk of adverse effects to river otter, spotted sandpiper, and osprey populations and to bald eagles. These latter risks are lower than the risk to the mink population. Further risk management recommendations regarding dioxins/furans are provided in Section 12.1.3.

12.1.2.2 COPCs Not Recommended as COCs

Inorganic Metals

None of the seven metals with $HQ \geq 1$ is recommended as a COC for assessing potential remedy protectiveness of ecological receptors. The rationales for exclusion are as follows:

- Aluminum poses potentially unacceptable risk only for mink. For the following reasons, it is not recommended as a COC:
 - Aluminum exceeds the dietary TRV only based on sediment ingestion, no prey samples exceed the effects threshold.
 - TRV is based on exposure of mice to a highly soluble ionic form of aluminum with higher bioavailability than typically found in the diet or drinking water.
 - Study Area sediment and surface water concentrations are similar to background.
- Antimony poses potentially unacceptable risk based only on the tissue-residue LOE for smallmouth bass. For the following reasons, it is not recommended as a COC:
 - Low frequency of TRV exceedance (1 of 32 [3.1%] smallmouth bass samples)

- Weakness of the exposure estimate (the single composite sample is an outlier for both antimony and lead, suggesting that a fish in the sample might have swallowed a fishing sinker)¹³
- Weakness of the effects estimate (TRV is based on a single study with a generic ACR applied)
- Absence of relationship between concentrations in sediment and co-located tissue samples (Windward 2009b)
- Discordance between the weaker tissue-residue LOE and the stronger surface water LOE (surface water TRV based on numerous exposure data and moderately sized Tier II effects dataset). Arsenic poses potentially unacceptable risk to benthic invertebrates based only on the tissue-residue LOE. It is not recommended as a COC for two reasons:
 - Low frequency of exceedance of the TRV (2 of 35 samples)
 - Low magnitude of the exceedance (maximum HQ = 1.5)
- Cadmium poses potentially unacceptable risk based only on the dietary LOE for juvenile Chinook salmon and sculpin. For the following reasons, it is not recommended as a COC:
 - Low frequency of TRV exceedance in sculpin prey samples (9 of 111 [8.1%] prey samples, with maximum HQ = 2.2; and 1 of 1,348 [$< 0.1\%$] sediment samples)
 - Weakness of the Chinook exposure estimate (juvenile Chinook were conservatively presumed to feed predominantly on benthic organisms; this feeding strategy is contrary to the literature, which shows they feed predominantly on pelagic organisms)
 - Uncertainty about the toxicological effects associated with the TRV (rockfish LOAEL setting the TRV is 2 to 3 orders of magnitude below the nine NOAELs from other studies, including four NOAELs and two LOAELs for salmonids)
 - Low magnitude of juvenile Chinook salmon dietary HQ (3.5 assuming mixed prey diet) when taking into account the likelihood that both exposure and effects are overestimated (per the two previous items)
 - Discordance of the dietary LOE with the surface water and tissue-residue LOEs (the cadmium AWQC is based on a very large dataset so is the strongest LOE; the tissue-residue LOE is weak because fish sequester or otherwise bioregulate inorganic metals)

¹³ Antimony can be mixed with lead as a hardener for lead-based products (ATSDR 1992). For example, one fish tackle supplier notes that fishing sinkers contain 94% lead and 6% antimony for hardness and color (Blue Ocean Tackle 2011).

- Copper poses potentially unacceptable risk based on the fish tissue-residue, fish dietary, sandpiper dietary, and the benthic invertebrate tissue-residue LOEs. For the following reasons, copper is not recommended as a fish COC:
 - Weakness of the tissue-residue LOE for inorganic metals (fish can actively bioregulate copper tissue concentrations; invertebrates sequester copper and in the case of crayfish, copper forms the basis of their hemoglobin)
 - Irreproducible toxicological effects associated with the dietary TRV (selected LOAEL could not be replicated in subsequent tests with the same species)
 - Selected LOAEL is barely above range of nutritional requirements found in the literature for some fish species
 - Discordance of the tissue and dietary LOEs with the stronger water LOE (which is based on numerous exposure data and a very large AWQC dataset showing that fish are not among the most sensitive species; absence of $HQ \geq 1$ via the water LOE is the strongest evidence for drawing risk conclusions)
 - Similarity of fish tissue concentrations in the Study Area and upriver

For the following reasons, copper is not recommended as a shorebird COC:

- Unlikely ecological significance of prey organism TRV exceedance (tissue-residue $HQ \geq 1$ in only one prey item, laboratory-exposed worms; $HQs < 1$ for a mixed-species diet).
- The selected TRV is less than the lowest bounded literature-reported NOAEL for birds.
- Low magnitude of TRV exceedance (maximum $HQ = 1.3$) considering the likely overestimates of exposure and effects (per the two previous items)

For the following reasons, copper is not recommended as a benthic invertebrate COC:

- Low magnitude of TRV exceedance (maximum $HQ = 2.6$)
- Weakness of the tissue-residue LOE for inorganic metals (invertebrates sequester copper and in the case of crayfish, copper forms the basis of their hemoglobin)
- Lead poses potentially unacceptable risk based on the tissue-residue LOE for peamouth and smallmouth bass, and on the dietary LOE for osprey and mink. It is not recommended as a fish COC for the following reasons:
 - Low frequency of tissue TRV exceedance (2 of 32 [6.2%] smallmouth bass and 1 of 4 [25%] peamouth samples)
 - Weakness of the exposure estimate (smallmouth bass concentration yielding high HQ [280] is an outlier for both antimony and lead in the same sample,

suggesting that a fish in the composite sample might have swallowed a fishing sinker)

- Discordance of tissue-residue LOE with dietary and water LOEs (based on a very large dataset, the lead AWQC is the strongest LOE; the tissue-residue LOE is weak because fish generally can sequester or otherwise bioregulate inorganic metals; the dietary LOE is more likely to overpredict than underpredict risk)

Lead is not recommended as a bird or mammal COC because the only sample yielding an $HQ \geq 1$ is the same outlier smallmouth bass sample as identified for antimony above

- Zinc poses potentially unacceptable risk for fish (sculpin, bass, pikeminnow), amphibians, and aquatic plants based only on the surface water LOE. It poses a potentially unacceptable risk to benthic invertebrates based on the surface water and tissue-residue LOEs. It is not recommended as a COC for the following reasons:
 - Low frequency of surface water TRV exceedance for all receptors (1 of 167 samples [$< 1\%$], with maximum $HQ = 1.2$)
 - Discordance of the stronger surface water LOE with the weaker tissue-residue and dietary LOEs for fish (surface water toxicity data were sufficient to derive AWQC; tissue-residue LOE is weak because fish generally can sequester or otherwise bioregulate inorganic metals; the dietary LOE is relatively weak because the TRV is based on only two studies)
 - The tissue-residue LOE for benthic invertebrates is a weak LOE

Organometals

- Mercury poses potentially unacceptable risk based on the dietary LOE for sculpin and bald eagle. It is not recommended as a fish COC because the dietary TRV was exceeded in only 1 of 1,345 sediment samples ($< 0.001\%$) and in no tissue samples. Mercury is not recommended as an eagle COC for the following reasons:
 - Discordance between the dietary and tissue-residue LOEs
 - Possible overestimate of bald eagle exposure when using osprey exposure as a surrogate because of greater proportion of terrestrial prey in the bald eagle diet
 - Low HQ (maximum $HQ = 1.7$) given the discordant LOEs and possibility that exposure is overestimated (per the previous two items)
 - Higher concentrations in upriver fish tissue than in Study Area fish tissue
- Monobutyltin poses potentially unacceptable risk based on the surface water LOE. It is not recommended as a COC for three reasons:

- Low frequency of surface water TRV exceedance (1 of 167 samples [$< 1\%$])
- Likely overestimate of toxicological effects associated with the TRV (which is based on the more toxic TBT)
- Low magnitude of exceedance ($HQ = 1.2$) considering the likely overestimate of effects and limited spatial extent of $HQ \geq 1$ (per the previous two items)
- TBT poses potentially unacceptable risk based on the dietary LOE for sculpin and tissue-residue LOE for benthic invertebrates. It is not recommended as a COC for fish for the following reasons:
 - Single dietary TRV exceedance (based on 1 lab worm sample of 81 prey samples [1.2%] and only when combined with sediment ingestion)
 - Low magnitude of exceedance (maximum $HQ = 1.0$)
 - Uncertainty about toxicological effects associated with the TRV (reproduction success was reduced at the TRV, but not dose-responsive)
 - Discordance of dietary LOE with the tissue-residue and water LOEs (TBT tissue residue is noted to be reliable predictor of toxicity and is the strongest LOE (Meador et al. 2002a))

It is not recommended as a COC for benthic invertebrates because of the following:

- The TRV was exceeded in empirical bioaccumulation samples only at one location.
- While predicted tissue residues exceeded the TRV more frequently, the moderate strength of the regression was highly influenced by the one high value in the dataset. The predicted tissue residues are uncertain and not supported by empirical data.
- The TRV is uncertain due to the inclusion of imposex—the endpoint that defined the lower distribution of the SSD, which set the TRV

PAHs¹⁴

Benzo(a)anthracene, benzo(a)pyrene, and naphthalene pose potentially unacceptable risk to benthic invertebrates, fish, amphibians, and aquatic plants based on the surface water LOE. Benzo(a)pyrene poses potentially unacceptable risk to spotted sandpiper based on the dietary LOE. None of these three individual PAHs is recommended as a COC for assessing potential remedy protectiveness of ecological receptors.¹⁵

¹⁴ Risk management recommendations regarding PAHs as they relate to risks from the TZW LOE and benthic AOCs are discussed separately in Sections 12.2 and 12.3, respectively.

¹⁵ In the TZW LOE, however, concordance of surface water and TZW exceedances at RM 6.4 to RM 6.5 supports identification of benzo(a)anthracene, benzo(a)pyrene and naphthalene as COCs for this location (see Section 12.2).

- Benzo(a)anthracene is not recommended as a COC for two reasons:
 - Low frequency of surface water TRV exceedance (2 of 245 samples [$< 1\%$], both between RM 6.4 and RM 6.5)¹⁶
 - Discordance of surface water LOE with dietary LOE for fish (benzo(a)anthracene did not screen in as a fish COPC by the dietary LOE)
- Benzo(a)pyrene is not recommended as a COC based on the surface water LOE for two reasons:
 - Low frequency of surface water TRV exceedance (3 of 122 [2.4%] near-bottom surface water samples, all from RM 6.4 to RM 6.5)¹⁷
 - Discordance of the surface water LOE with the dietary LOE for fish (benzo(a)pyrene did not screen in as a fish COPC by the dietary LOE)
- Benzo(a)pyrene is not recommended as a COC based on the bird dietary LOE for two reasons:
 - Low frequency of dietary TRV exceedance for spotted sandpiper (1 of 27 [3.7%] lab worm samples assuming lab worm-only diet; all HQs < 1 for clam-only diet)
 - Low magnitude of exceedance (maximum HQ = 1.6) considering potential overestimate of exposure by presuming lab worm-only diet
- Naphthalene is not recommended as a COC for two reasons:
 - Low frequency of surface water TRV exceedance (10 of 268 [3.7%] samples, all from west side of RM 6.4 to RM 6.5 during a single sampling event [the May 2005 non-LWG sampling event])¹⁸
 - Discordance of the surface water LOE with the dietary LOE for fish (naphthalene did not screen in as a fish COPC by the dietary LOE)

Phthalates

Neither of the two phthalates is recommended as a COC:

- BEHP poses potentially unacceptable risk based on the benthic invertebrate and fish tissue-residue and surface water LOEs. It is not recommended as a COC for several reasons:

¹⁶ In the TZW LOE, however, concordance of surface water and TZW exceedances at this sampling location support identification of benzo(a)anthracene as a COC for this location (see Section 12.2).

¹⁷ In the TZW LOE, however, concordance of surface water and TZW exceedances at this sampling location support identification of benzo(a)pyrene as a COC for this location (see Section 12.2).

¹⁸ In the TZW LOE, however, concordance of surface water and TZW exceedances at this sampling location support identification of naphthalene as a COC for this location (see Section 12.2).

- Low frequency of surface water TRV exceedance (2 of 190 samples [1.1%])
- Low frequency of fish tissue-residue TRV exceedance (1 of 38 sculpin samples [2.6%], 2 of 32 smallmouth bass samples [6.3%]) and low frequency of the benthic invertebrate tissue-residue TRV exceedance (1 of 35 clam samples or 3%)
- Low magnitude of exceedance for fish tissue TRV (maximum HQ = 2.9) and for benthic invertebrate TRV (maximum HQ = 2.8)
- Absence of toxicological effects associated with the tissue TRV (which is based on an unbounded NOAEL)
- Absence of relationship between concentrations in co-located sediment and tissue samples
- Dibutyl phthalate poses potentially unacceptable risk based on the dietary LOE for spotted sandpiper. It is not recommended as a COC for several reasons:
 - Low frequency of dietary TRV exceedance (1 of 28 clam samples [3.6%], no worm samples)
 - Low magnitude of dietary TRV exceedance (maximum HQ = 1.4 for clam-only diet; maximum HQ < 1 for worm-only diet)
 - Absence of a relationship between concentrations in co-located sediment and tissue samples
 - Higher sediment concentrations in background than in Study Area

Pesticides

None of the three organochlorine pesticides is recommended as a COC for assessing potential remedy protectiveness of ecological receptors:

- Aldrin poses potentially unacceptable risk based on the dietary LOE for spotted sandpiper. It is not recommended as a COC for two reasons:
 - Low frequency of dietary TRV exceedance (1 of 27 lab worm samples [3.7%])
 - Low magnitude of exceedance (maximum HQ = 1.4 based on the only lab worm sample that yields an HQ ≥ 1 ; HQ < 1 for clam-only and mixed diets)
- Total DDX poses potentially unacceptable risk based on the tissue-residue LOE for sculpin and benthic invertebrates; the dietary LOE for spotted sandpiper; the egg LOE for bald eagle; and the surface water LOE for the benthic community, sculpin, amphibians, and aquatic plants. The rationale for exclusion from the list

of recommended COCs varies with LOE.¹⁹ DDx is not recommended as a COC for the following reasons:

- Low frequency of TRV exceedance (1 of 170 samples [$<1\%$]) in surface water based on N-qualified data, indicating interference from another analyte
- Low frequency of exceedance in empirical benthic tissue residue (2 of 35 worm samples or 6%)
- Low frequency of exceedance in predicted benthic tissue residues (up to 15 samples of 1,128 or 1.3%) and approximately half of which are based on N-qualified data
- Low frequency of TRV exceedance (2 of 27 lab worm samples [7.4%]) used in the dietary LOE for sandpiper
- Low magnitude of exceedance of TRV for sandpiper diet (maximum HQ = 1.4 assuming lab worm-only diet; HQ < 1 for all clam-only and mixed diets)
- Questionable relevance of estimated exposure for the bird egg LOE for bald eagle (there is significant uncertainty about the source of DDx residues in the osprey eggs collected from the Study Area because the adults overwinter in Mexico and Central America, nesting and laying eggs shortly after returning to the lower Willamette (Henny et al. 2004))
- Potential risk of adverse effects on bald eagles is present because NOAEL HQs are ≥ 1 in eggs from two of five exposure areas; because both were below the LOAEL, there is no empirical evidence of potential risk.
- All egg total DDx concentrations were below the recommended effects threshold reported in Elliott and Harris (2001\2002) based on a comprehensive review of the available bald eagle toxicological effects data
- Absence of relationship between concentrations in osprey egg samples and nearby sediment (NOAEL HQ ≥ 1 in eggs from two of five exposure areas, but NOAEL HQ < 1 in eggs from where sediment DDx concentrations were highest)
- Discordance of LOEs (mixed species dietary NOAEL HQs < 1 in all exposure areas)
- 4,4'-DDD poses potentially unacceptable risk based on the tissue-residue LOE for benthic invertebrates. This contaminant is not recommended as a COC for the following reasons:
 - Low frequency of TRV exceedance (1 of 35 samples or $< 3\%$)

¹⁹ Total DDx and 4,4'-DDT are recommended as TZW COCs in the TZW sampling area at ~ RM 7.4W (see Section 12.2).

- Low magnitude of the exceedance (HQ = 1.2)

VOCs

Two VOCs (ethylbenzene and trichloroethene) measured in surface water exceeded their respective TRVs; however, neither is recommended as a COC based on the following rationale:

- Low frequency of exceedance; TRV exceeded in 1 of 23 (4%) samples collected from ~ RM 6.5 (west bank) during one sampling event
- Low magnitude of exceedance of the TRV for ethylbenzene (HQ = 1.6)

12.1.3 Risk Management Recommendations for Recommended COCs

Based on the information presented in Section 12.1, total PCBs and total TEQ pose the primary risks to fish and wildlife. The remainder of this section provides additional risk management recommendations for these recommended COCs:

- Section 12.1.3.1 recommends the use of mink to evaluate total PCB and total TEQ remedies.
- Section 12.1.3.2 examines relationship between PCB and TEQ risk.
- Section 12.1.3.3 discusses potential problems with the use of the bird egg LOE as an evaluation tool for potential remedies.

12.1.3.1 Receptors of Concern for Purposes of Assessing the Protectiveness of Potential Remedies in the FS Analysis of Alternatives

Total PCBs is recommended as a COC because exposure poses a risk of ecologically significant adverse effects to mink and river otter populations. Total PCBs also poses lower risk of ecologically significant adverse effects to benthic invertebrates, spotted sandpiper, osprey, sculpin, and smallmouth bass populations and to bald eagles. Total TEQ is recommended as a COC because exposure poses a risk of ecologically significant adverse effects to mink populations. Total TEQ also poses lower risk of adverse effects to river otter, spotted sandpiper, and osprey populations and to bald eagles.

For the dietary LOE, HQs are a function of food and sediment ingestion rates relative to the organism's body weight, the COPC concentrations in prey and sediment, and the TRV. Of the receptors at risk from PCBs and total TEQ via the dietary LOE, mink has the lowest TRVs. The bird PCB LOAEL TRV is higher than that of mink by a factor of 16, and the bird total TEQ LOAEL TRV is higher than that of mink by a factor of 64, indicating that risk to mink occurs at lower dietary doses.

Given the same sediment and prey data, dietary risk estimates for mink will always be higher and more widespread than those for the other receptors. Food and sediment ingestion rates as a function of body weight are higher for mink than for otter; and they are higher for birds than for mink (by a factor ranging from 1.3 for osprey to 7 for spotted sandpiper). However, the difference in TRVs (for both total PCBs and total TEQ) more

than offsets the difference in ingestion rates. Although a mink population is not known to be present in the Study Area, mink are assumed to forage in all areas of the Study Area and to prey on small- and large-home-range fish. Analysis of remedial alternatives for mink will thus be protective of other receptors in the Study Area potentially affected by PCBs and dioxins.

Predicted mink risk is based on species-specific effects data, making mink risk predictions a relatively strong basis for risk management decisions. This is not the case for the other receptors (predicted risks are not based on species-specific effects data), whose conclusions therefore provide a less certain basis for risk management recommendations. Because the available data suggest that mink are quite sensitive to PCBs and dioxins/furans, and probably more so than the other receptors at risk, the mink population should be the receptor of concern when assessing ecological risk reduction for the remedial alternatives (for total PCBs and total TEQ).

Because protection of other receptors by mink is contingent on the habitat use, prey, and home-range assumptions used for the BERA, any alteration of these assumptions for analysis of uncertainties in the FS should be examined to ensure that protection of all receptors at risk from PCBs and TEQ are still protected under alternate assumptions for mink.

Because the relationship between sediment contamination and bird egg tissue concentrations is highly uncertain, the tissue-residue LOE has limited utility as a tool for assessing the protectiveness of potential remedies in the FS analysis of alternatives. This is discussed further in Section 12.1.3.3.

12.1.3.2 Relationship Between PCB and TEQ Risk

Total TEQ is the sum of multiple PCB and dioxin/furan congeners, each weighted by their toxicity relative to that of the most toxic congener (2,3,7,8-TCDD). TEQ concentrations for birds and mammals were calculated as the sum of individual PCB and dioxin/furan congener concentrations weighted by their TEFs. The PCB TEQ is the TEF-weighted sum of only the dioxin-like PCB congener concentrations, the total dioxin/furan TEQ is the TEF-weighted sum of only the dioxin/furan congener concentrations, and total TEQ is the sum of the PCB TEQ and the total dioxin/furan TEQ. TEF values for a given congener generally fall within a range of about an order of magnitude for mammals (Sanderson and Van den Berg 1999); TEFs for birds are more uncertain (Van den Berg et al. 1998). Because of this uncertainty, TEQ risks may be over- or underestimated.

As with total PCBs, mink is the receptor most sensitive to dioxins/furans and subject to the greatest spatial extent of TEQ risk in the Study Area. PCBs are responsible for the majority of total TEQ risk, in that PCB TEQ HQs generally constitute the majority of the total TEQ HQs. For example, of the 15 (out of 109) potential prey samples with mink total TEQ HQ ≥ 1 , 7 exceed the TRV for PCB TEQ but only 4 exceed the TRV for total dioxin/furan TEQ (see Attachment 17). No individual samples result in an exceedance of both the PCB TEQ TRV and the dioxin/furan TEQ.

Because total TEQ risk is largely driven by PCB, and redundant with total PCB risk (with the four exceptions noted above), and because adverse effects in mink are better correlated with total PCB exposures than with TEQ exposures (Fuchsman et al. 2007), the FS analysis of alternatives should focus primarily, but not exclusively, on evaluating whether remedies protect the mink population from risk due to exposure to total PCBs.

12.1.3.3 Bird Egg LOE and the FS

PCBs and total TEQ pose low risk to birds based on the tissue-residue LOE. It is recommended that the bird egg LOE not be used to develop and evaluate remedial alternatives in the FS. Risk to osprey and bald eagle based on the egg LOE cannot be directly compared with dietary risks. Egg tissue concentrations might reflect exposure to contaminated prey from the Study Area. Alternatively, inasmuch as osprey lay eggs shortly after returning to the Study Area from overwintering in Mexico and Central America, the egg residues might reflect exposure to contaminants outside of the Study Area. Furthermore, the bioaccumulation relationship from prey to egg is not well-characterized, rendering predictions based on this relationship highly uncertain.

A statistical evaluation was conducted to determine if a relationship between fish tissue and bird egg tissue concentrations in the Study Area could be expressed using biomagnification regressions (BMRs). A BMR expresses the relationship between fish prey and bird egg tissue concentrations based on co-located data rather than based on an average ratio. BMRs were calculated based on the method by Burkhard (2009) using co-located (within 1 mile) composite fish tissue and egg concentrations from seven locations throughout the Willamette River (Henny et al. 2003; 2009). Several possible linear tissue-sediment models were screened. No significant relationship (i.e., no BMR) could be found for any bird egg COPC based on the criteria of a significantly positive slope at a $p = 0.05$ and an $r^2 > 0.030$, except total TEQ ($r^2 = 0.52$). For total TEQ, application of the BMR to the Study Area requires extrapolation outside of the dataset, thus rendering the relationship uncertain. The implication is that the available dataset is insufficient to estimate a reliable BMR.

Because mink is the receptor most sensitive to PCBs and dioxins/furans, it is recommended that from an ecological risk management perspective, FS analyses should focus primarily on the mink dietary risk reduction associated with the remedial alternatives.

12.2 TZW RISK MANAGEMENT RECOMMENDATIONS

The TZW LOE was used to assess risks to benthic invertebrate, benthic fish (i.e., sculpin and lamprey ammocoetes), aquatic plant, and amphibian populations and communities. Pacific lamprey are identified in EPA's Problem Formulation as a "species of special concern" with direction to assess risk at the organism level. Measured TZW concentrations exceed water TRVs in all of the TZW sampling areas; by EPA's direction individual lamprey ammocoetes are exposed to potentially unacceptable risk. The degree to which TZW poses potentially unacceptable risk to individual lamprey ammocoetes is uncertain. Lamprey ammocoete toxicity testing has demonstrated their relative

insensitivity to toxicants across six modes of action (Andersen et al. 2010). It is probable that the BERA overestimates both lamprey ammocoete exposure and effects, to an unquantified degree.

The TZW samples evaluated in this assessment were collected primarily during a 2005 sampling effort focused offshore of nine²⁰ upland sites with known or likely pathways for discharge of upland contaminated groundwater. The primary objective of the RI groundwater pathway assessment was to evaluate whether transport pathways from upland contaminated groundwater plumes to the river were complete. Therefore, TZW target analyte lists varied from site to site and were derived primarily based on the COIs in the upland contaminated groundwater plumes. Consequently, not all COIs in sediments were analyzed in TZW samples. As described in Sections 4.4.3.1 and 6.1.5.2 of the draft final RI (Integral et al. 2011), there also might be other groundwater plumes in the Study Area discharging into river sediments where TZW samples have not been collected.

TZW sampling focused on sites with contaminated groundwater pathways that were a potential concern. Where these groundwater pathways are confirmed to be a concern, they will be addressed through source control. Source controls should be in place prior to implementation of sediment remedies, particularly those associated with upland sources (EPA 2002b, 2005a) in order to prevent recontamination. These source control actions will reduce contaminant flux to the river and accelerate recovery. Source controls will reduce baseline risk by intercepting ongoing contaminant migration. While the residual contaminated groundwater plumes may remain near the mudline, they will attenuate over time. Because source controls should precede the sediment remedy, the magnitude of potential risk identified in the BERA should be diminished when the sediment remedy is implemented.

The TZW LOE was evaluated by comparing TZW COPC concentrations in individual samples to water effect thresholds. EPA directed the LWG to assume that benthic organisms would be exposed to undiluted shallow (0 to 38 cm) TZW, an assumption that the LWG found to be highly conservative. As discussed in Section 6.6.3.3, actual TZW exposure is probably much lower because of feeding habits, burrowing behavior, avoidance of low oxygen levels at the TZW sample depths, and low food content in sediments at the TZW sample depths.

It is recommended that only those TZW COPCs with $HQ \geq 100$ be considered as COCs to develop and evaluate remedial alternatives that are protective of ecological resources.²¹ This recommendation is based on two factors. First, by definition any contaminant with $HQ \geq 1$ poses potentially unacceptable risk, but the evidence presented in Section 6.6.3.3

²⁰ The area offshore of the Arkema site was divided into two areas (the acid plant area and the chlorate plant area).

²¹ There is uncertainty associated with 4,4'-DDT and total DDx as COCs because HQs based on filtered samples are less than 100. This suggests that the risk from DDx compounds in TZW may be lower than indicated by the maximum concentrations in unfiltered samples due to lower bioavailability of the particulate bound fraction of the contaminant.

strongly supports the position that the potential for unacceptable risk at HQs < 10 is very small. Therefore, a factor of 10 was applied to account for the evidence that benthic receptors are not directly exposed to undiluted TZW. Second, EPA guidance (EPA 2005a) states that remedies should be evaluated under the assumption that sources of COPCs to the groundwater plume have been controlled. The effect of source control should be to reduce the potential flux of groundwater COPCs into the shallow transition zone prior to sediment remediation. An additional factor of 10 was applied to account for the control of COPC sources.

Almost all metals measured in TZW are common crustal elements. Barium, iron, and manganese are among the most common metals associated with sediments. These same metals are also associated with the highest HQs in the risk characterization, but there is substantial uncertainty that their source is ubiquitously anthropogenic. It is recommended that TZW concentrations of these metals not be used to assess remedy effectiveness.

Given the foregoing, TZW COC recommendations for each site are provided in Table 12-2.

Table 12-2. COC Recommendations for COPCs with HQs ≥ 100 at TZW Sampling Areas

COPC	Maximum HQ ≥ 100									
	ARCO	Arkema		Exxon Mobil	Gasco	Gunderson	Kinder Morgan	Rhône-Poulenc	Siltronic	Willbridge
		Acid Plant	Chlorate Plant							
Contaminants Recommended as TZW COCs										
Benzo(a)anthracene					120				1,200	
Benzo(a)pyrene					210				2,700	
Naphthalene					260				1,100	
4,4'-DDT		160 ^a								
Total DDx		280 ^a								
Chlorobenzene		190								
cis-1,2-Dichloroethene									110	
Trichloroethene									1,900	
Cyanide					4,400				130	
Carbon disulfide					870					
Contaminants Not Recommended as TZW COCs										
Barium (total)		610	1,100					170		
Iron (total)		110	250	110	130				180	120
Manganese (total)			550	150	130			130		110
Gasoline-range aliphatic hydrocarbons C10-C12 ^b					540				150	

^a Maximum HQs are based on unfiltered samples. Maximum HQs for filtered samples would be 2.8 for 4,4'-DDT (however, this contaminant was never detected) and 14.5 for total DDx.

^b Petroleum hydrocarbons may contribute to risks to ecological receptors; however, petroleum is not considered a CERCLA contaminant.

CERCLA – Comprehensive Environmental Response,
Compensation, and Liability Act

DDD – dichlorodiphenyldichloroethane
DDE – dichlorodiphenyldichloroethylene
DDT – dichlorodiphenyltrichloroethane

HQ – hazard quotient
total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-
DDD, 2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT)

COC – contaminant of concern

COPC – contaminant of potential concern

Potential remedies should be evaluated in the FS for the degree to which they protect benthic invertebrate communities and individual Pacific lamprey ammocoetes from risk due to contaminated groundwater discharge, assuming that groundwater source control measures have been implemented.

12.3 BENTHIC RISK MANAGEMENT RECOMMENDATIONS

The primary LOE for identifying benthic community risks is based on sediment toxicity (both measured and predicted based on multivariate statistical models [FPM and LRM]); however, the risk assessment methodologies are designed to address chemical mixtures. The results are correlative and do not conclusively identify contaminants causing toxicity.²² Contaminants whose sediment concentrations, when considered as a group (i.e., in aggregate), appear to help explain the observed toxicity based on the FPM and LRM are presented in Table 12-3.²³

Table 12-3. Contaminants Potentially Contributing to Benthic Risk Based on Predicted Sediment Toxicity LOE

Contaminant	
Metals	
Cadmium	Lead
Chromium ^b	Mercury ^b
Copper	Silver
PAHs	
2-Methylnaphthalene	Dibenzo(a,h)anthracene
Acenaphthene	Fluoranthene
Acenaphthylene	Fluorene
Anthracene	Indeno(1,2,3-cd)pyrene
Benzo(a)anthracene	Phenanthrene
Benzo(b)fluoranthene	Pyrene
Benzo(b+k)fluoranthene	Total HPAHs
Benzo(g,h,i)perylene	Total LPAHs ^b
Benzo(k)fluoranthene	Total PAHs

²² Risk conclusions based on the secondary benthic LOEs—tissue residue, surface water, and TZW—can identify COCs and are noted in Sections 12.1 and 12.2, where these LOEs support the identification of COCs.

²³ The contaminant list is a combination of SQVs derived using the FPM and the LRM. Each SQV has a different reporting basis depending on the normalization selected for the model. All FPM SQVs are dry-weight normalized. LRM SQVs used a number of different normalizations including dry-weight, organic carbon, percent fines and combinations of normalizations.

Table 12-3. Contaminants Potentially Contributing to Benthic Risk Based on Predicted Sediment Toxicity LOE

Contaminant	
Chrysene	
Phthalates	
Dibutyl phthalate	
SVOCs	
Benzyl alcohol	Dibenzofuran ^b
1,2-Dichlorobenzene	Carbazole ^b
Phenols	
4-Methylphenol ^a	Phenol
PCBs	
Total PCBs ^b	
Pesticides	
2,4'-DDD	beta-HCH
4,4'-DDD	delta-HCH ^b
4,4'-DDE	Dieldrin
4,4'-DDT	Endrin
Sum DDD ^b	Endrin ketone
Sum DDE	cis-Chlordane
Sum DDT	Total endosulfan ^b
Total DDx	
Petroleum Hydrocarbons	
Diesel-range hydrocarbons	

^a All SQVs derived from the FPM are less than the apparent effect threshold and therefore may contribute to false predictions of toxicity.

^b FPM SQVs based on one or two endpoints are less than the apparent effect threshold and may contribute to false predictions of toxicity

COPC – contaminant of potential concern

DDD – dichlorodiphenyldichloroethane

DDE – dichlorodiphenyldichloroethylene

DDT – dichlorodiphenyltrichloroethane

FPM – floating percentile model

HCH – hexachlorocyclohexane

HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

LOE – line of evidence

LPAH – low-molecular-weight polycyclic aromatic hydrocarbon

PAH – polycyclic aromatic hydrocarbon

PCB – polychlorinated biphenyl

SVOC – semivolatile organic compound

total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT)

Because the primary benthic LOE (bioassay results) does not identify the cause of the empirical toxicity (i.e., specific COPCs or other factors), the risk management recommendations focus on two other questions:

1. Where were potentially unacceptable benthic community risks occurring in the Study Area at the time of the BERA data collection?
2. What tools from the BERA can be used in the FS analysis of alternatives to assess the effectiveness of potential remedies on protecting the benthic community?

The remainder of this section is arranged around these two questions. Section 12.3.1 outlines the guidelines EPA provided about how to answer them. Section 12.3.2 answers the first question by presenting recommended benthic AOCs. Section 12.3.3 answers the second question by recommending tools by which to assess the effect of potential remedies on the benthic community in the FS analysis of alternatives.

12.3.1 EPA Guidelines for Evaluating Benthic Risk in the Feasibility Study

The LWG and EPA have been working on benthic risk management recommendations since early 2010, following guidelines EPA in an April 21, 2010 letter (EPA 2010a). The guidelines provide direction for evaluating benthic risk in the draft FS. Specifically, EPA described its primary goals for the FS analysis of alternatives for benthic assessment endpoints:

- Define areas that pose unacceptable risk to the benthic community
- Define the areas and volume of contamination that may pose risk to the benthic community
- Evaluate remedial action alternatives and effectiveness (did it meet the RAO)

The letter also provided guidelines for evaluating remedy effectiveness:

- All benthic SQGs in the March 24, 2010 list will be included in the analysis. If specific SQGs are found to be inconsistent with other LOEs listed below, EPA will review the analysis and determine whether these should be included in the draft FS.²⁴
- Sediment toxicity bioassays will form the primary LOE for this analysis. The sediment toxicity LOE will include level 2 (moderate) and level 3 (severe) effects for all endpoints (chironomus [sic] biomass and mortality and hyalella [sic] biomass and mortality).

²⁴ The SQVs have subsequently been revised based on additional modeling and negotiations between the LWG and EPA, as documented in item 11 of Attachment B to a January 12, 2011, LWG letter to EPA (LWG 2011a), the attachment to a February 25, 2011, RI/FS schedule letter from EPA to the LWG (Humphrey 2011), and the LWG's March 9, 2011, draft response (LWG 2011b) to EPA's February 25, 2011, letter.

- The analysis will consider the number and degree of exceedance of SQGs.
- The analysis will consider other LOEs such as TZW compared to ambient water quality criteria for the protection of aquatic life and benthic tissue TRVs.
- The analysis will consider the presence/absence of nearby sources and examine benthic community structure (e.g., via sediment profile imaging and related information).
- The analysis will consider data quality and data density issues for the SQGs.

The LWG's implementation of these guidelines is known by EPA and the LWG as the "comprehensive benthic approach." Developed by the LWG after receiving the EPA's April 21, 2010, directives and guidelines (EPA 2010a), the comprehensive benthic approach was first presented informally to EPA (Eric Blischke and Burt Shephard) by the LWG (John Toll and Jim McKenna) on July 20, 2010, to elicit early feedback. It was formally presented to EPA during the September 29, 2010, LWG Small Technical Group Benthic Toxicity AOPCs Meeting with EPA. Item 11 in Attachment B to the LWG's January 12, 2011, letter to EPA (LWG 2011a), and the attachment to EPA's February 25, 2011, response letter to the LWG (Humphrey 2011) document the decision to proceed with an updated version of the comprehensive benthic approach.

12.3.2 Recommended Benthic Areas of Concern for FS Evaluation

Recommended benthic AOCs, based on the LWG's application of the comprehensive benthic approach upon completion of the draft final BERA, are shown on Maps 12-1a and 12-1b. Sediment toxicity bioassays form the primary LOE for the comprehensive benthic approach used to delineate the recommended benthic AOCs, as per the EPA April 21, 2010, guidelines (EPA 2010a). Predicted toxicity (based on multiple sets of SQVs) and tissue residues (both empirical and predicted) provide secondary LOEs to identify benthic risk areas. TZW and surface water were used as supporting LOEs.

SPI data were not used in the development of AOCs because the sampling program was not designed to link SPI image locations with toxicity sampling locations and in turn allow an assessment of the relationship between benthic community successional stage and contaminant effects. Details of the approach used to identify recommended benthic AOCs are as follows:

- Locations with empirical bioassay results indicating significant toxicity were identified.
 - One toxicity endpoint (*Chironomus* biomass or growth, *Hyalella* biomass or growth) exceeding an L3 threshold or two endpoints exceeding an L2 endpoint were considered significant toxicity.

- Locations where significant sediment toxicity is predicted based on sediment chemistry exceeding an MQ of 0.7 or a pMax of 0.59 were identified.
 - Sampling locations where both the MQ and the pMax thresholds were exceeded were considered toxic.
 - Sampling locations where neither the MQ or pMax threshold was exceeded were considered non-toxic.
 - Sampling locations where the models disagreed (i.e., either the MQ or the pMax threshold was exceeded, but not both) were considered uncertain.
- Locations where empirical tissue residues or, in the absence of empirical tissue residue data, predicted tissue residues exceeded their TRVs were identified.
 - The evidence of risk provided by measured or predicted exceedance of metals TRVs was considered weak because of species-specific differences in metals sequestration or other bioregulation.
 - The evidence of risk provided by predicted exceedance of the TBT TRV was considered weak because of high uncertainty in the TBT bioaccumulation model.
- TZW exceedance areas with HQs > 100 were delineated.
- All LOEs were overlaid on a map.
 - Areas where two or more adjacent empirical bioassay sampling locations indicate significant toxicity were identified as benthic AOCs.
 - Areas where risks were identified at two or more adjacent sampling locations based on chemistry LOEs (predicted toxicity, empirical or predicted bioaccumulation) or a combination of bioassay and chemistry LOEs were identified as benthic AOCs.
 - TZW exceedance areas were identified as benthic AOCs.
- Boundaries of the benthic AOCs split the distance between sampling locations exceeding criteria and surrounding clean sampling locations except where:
 - Other physical features were present (e.g., pier, channel edge, property boundary), in which case the boundary was drawn at the physical features.
 - The nearest sampling location was at a distance greater than 200 ft, in which case the boundary was drawn at a subjective distance less than halfway to nearest sampling location.

12.3.3 Benthic Assessment Tools for the FS Analysis of Alternatives

Bioassays cannot form the primary LOE for the FS analysis of alternatives, because the analysis is of potential future conditions. Therefore, the sediment chemistry LOE, as applied in the comprehensive benthic approach, will have to be used to judge protectiveness of potential remedies. The comprehensive benthic approach uses

concordance between an MQ based on the site-specific SQVs and the predicted pMax to identify benthic risk areas. EPA selected the MQ threshold of 0.7 and the pMax threshold of 0.59 that the LWG used in defining benthic AOCs. These same thresholds should be used to evaluate the protectiveness of potential remedies. The analysis of alternatives should also consider whether and how much natural recovery would occur prior to implementing active remedies. Per EPA guidance (EPA 2002b, 2005a), the analysis should presume that source control measures will be in place.

12.4 SUMMARY OF RISK MANAGEMENT RECOMMENDATIONS

The purpose of the risk management recommendations provided in Section 12 is to identify COCs, receptors, and AOCs that the LWG considers necessary and sufficient to develop and evaluate remedial alternatives that are protective of ecological resources. The FS will also evaluate whether remedial alternatives for these COCs, receptor and AOCs address the full list of contaminants potentially posing unacceptable risk.

In summary, the following are recommended as receptor-COC pairs of concern for further consideration in the FS:

- For non-benthic invertebrate receptors, total PCBs and total TEQ are the recommended COCs. Mink is the recommended receptor of concern. Most of the contaminants posing potentially unacceptable risk were not recommended as COCs for the non-benthic receptors based on risk characterization considerations (magnitude, spatial extent, and ecological significance of HQs ≥ 1). This list includes all the metals, butyltin, phthalate, pesticide, and VOC COPCs.
- For aquatic receptors exposed via TZW, 4,4'-DDT, total DDx,²⁵ chlorobenzene, benzo(a)anthracene, benzo(a)pyrene, naphthalene, carbon disulfide, cyanide, cis-1,2-dichloroethene, and trichloroethene are the recommended COCs. These recommendations presume that contaminated groundwater source control measures will be implemented prior to sediment remedies. ODEQ is working with upland property owners to implement contaminated groundwater source control measures prior to sediment remedies.
- For benthic organisms, recommended benthic AOCs were mapped by applying the comprehensive benthic approach based on EPA's April 21, 2010, guidelines for assessing benthic risk in the FS (EPA 2010a). The FS analysis of alternatives will have to rely on the predicted toxicity metrics to evaluate potential remedies and should take into account sediment quality changes that will take place before active implementation of remedies.

²⁵ There is uncertainty associated with 4,4'-DDT and total DDx as COCs because HQs based on filtered samples are less than 100. This suggests that the risk from DDx compounds in TZW may be lower than indicated by the maximum concentrations in unfiltered samples because of the lower bioavailability of the particulate-bound fraction of the contaminant.

13.0 REFERENCES

Adams SM, Ham KD, Greeley MS, LeHew RF, Hinton DE, Saylor CF. 1996. Downstream gradients in bioindicator responses: point source contaminant effects on fish health. *Can J Fish Aquat Sci* 53:2177-2187.

Adolfson, Walker Macy, Esther Lev, Winterowd, Ecotrust. 2000. Willamette River inventory: Natural resources. Public review draft. Prepared for Bureau of Planning, City of Portland. August 2000. Adolfson Associates Inc., Walker Macy, Esther Lev Environmental Consulting, Winterowd Planning Services, and Ecotrust, Portland, OR.

Allen AW. 1986. Habitat suitability index models: mink, revised. *Biol rep* 82(10.127). US Fish and Wildlife Service, Washington, DC.

Andersen HB, Caldwell RS, Toll J, Do T, Saban L. 2010. Sensitivity of lamprey ammocoetes to six chemicals. *Arch Environ Contam Toxicol* 59:622-631.

Anderson MR. 2008. Description and use of the RSET floating percentile method spreadsheets. Draft. September 6, 2008. Regional Sediment Evaluation Team, Oregon Department of Environmental Quality, Portland, OR.

Anthony RG, Garrett MG, Schuler CA. 1993. Environmental contaminants in bald eagles in the Columbia River estuary. *J Wildl Manage* 57(1):10-19.

Arnot JA, Gobas FAPC. 2004. A food web bioaccumulation model for organic chemicals in aquatic ecosystems. *Environ Toxicol Chem* 23:2343-2355.

ASTM. 2007. Standard test method for measuring the toxicity of sediment-associated contaminants with freshwater invertebrates. Method E1706-00. Annual book of ASTM standards, vol 11.06 [online]. American Society for Testing and Materials, West Conshohocken, PA. Available from: http://www.astm.org/cgi-bin/SoftCart.exe/DATABASE.CART/REDLINE_PAGES/E1706.htm?L+mystore+xzea3685+1187379245.

Atchison GJ, Henry MG, Sandheinrich MB. 1987. Effects of metals on fish behavior: a review. *Environ Biol Fish* 18(1):11-25.

ATSDR. 1992. Toxicological profile for antimony and compounds. Agency for Toxic Substances and Disease Registry, Atlanta, GA.

ATSDR. 2008. Toxicological profile for aluminum. September 2008. Agency for Toxic Substances and Disease Registry, Atlanta, GA.

Aulerich RJ, Ringer RK. 1977. Current status of PCB toxicity to mink, and effect on their reproduction. *Arch Environ Contam Toxicol* 6:279-292.

Aulerich RJ, Bursian SJ, Breslin WJ, Olson BA, Ringer RK. 1985. Toxicological manifestations of 2,4,5,2',4',5'-, 2,3,6,2',3',6'-, and 3,4,5,3',4',5'- hexachlorobiphenyl and Aroclor 1254 in mink. *J Toxicol Environ Health* 5:63-79.

Avocet. 2003. Development of freshwater sediment quality values for use in Washington State. Phase II report: Development and recommendation of SQVs for freshwater sediments in Washington State. Publication No. 03-09-088. Prepared for Washington Department of Ecology. Avocet Consulting, Kenmore, WA.

Baldigo BP, Baudanza TP. 2001. Avoidance response and mortality of juvenile brown trout (*Salmo trutta*) in tests with copper-sulfate-treated water from West Branch Reservoir, Putnam County, New York. USGS water resources investigations report 99-4237. US Geological Survey, Troy, NY.

Baldwin DH, Sandahl JF, Labenia JS, Scholz NL. 2003. Sublethal effects of copper on coho salmon: impacts on nonoverlapping receptor pathways in the peripheral olfactory nervous system. *Environ Toxicol Chem* 22(10):2266-2274.

Bartlett L, Rabe FW, Funk WH. 1974. Effects of copper, zinc and cadmium on *Selanastrum capricornutum*. *Wat Res* 8:179-185.

Baumann PC. 2000. Health of bullhead in an urban fishery after remedial dredging. Final report, January 31, 2000. Prepared for Great Lakes National Program Office, US Environmental Protection Agency [online]. US Geological Survey Field Research Station, Ohio State University, Columbus, OH. [Cited 10/30/2006.] Available from: <http://epa.gov/glnpo/sediment/Bullhead/index.html>.

Baumann PC, Harshbarger JC. 1995. Decline in liver neoplasms in wild brown bullhead catfish after coking plant closes and environmental PAHs plummet. *Environ Health Perspect* 103(2):168-170.

Baumann PC, Smith WD, Ribick M. 1982. Polynuclear aromatic hydrocarbons (PAH) residue and hepatic tumor incidence in two populations of brown bullhead (*Ictalurus nebulosus*). In: Cook M, Dennis AJ, Fisher G, eds, Polynuclear aromatic hydrocarbons: physical and biological chemistry. Battelle Press, Columbus, OH, pp 93-102.

Baumann PC, Smith WD, Parland WK. 1987. Tumor frequencies and contaminant concentrations in brown bullheads from an industrialized river and a recreational lake. *Trans Am Fish Soc* 116:79-86.

Baumann PC, Smith IR, Metcalfe CD. 1996. Linkages between chemical contaminants and tumors in benthic Great Lakes fish. *J Great Lakes Res* 22(2):131-152.

Beak. 2000. Ross Island and lower Willamette River fish monitoring study - 1999. Prepared for Ross Island Sand & Gravel, Landau Associates, and Perkins Coie, LLP. Beak Consultants, Inc., Portland, OR.

Beamesderfer RCP, J.C. Elliott JC, C.A. Foster CA. 1998. Methods for ageing Columbia River white sturgeon from pectoral fin rays. Report A. 1) Description of the life history and population dynamics of subadult and adult white sturgeon in the Columbia River between Bonneville and McNary dams, and 2) Evaluation of the need and identification of potential methods for protecting, mitigating, and enhancing white sturgeon populations in the Columbia River downstream from McNary Dam. In: Nigro AA, ed, Status and habitat requirements of white sturgeon populations in the Columbia River downstream from McNary Dam. Annual progress report to Bonneville Power Administration, Portland, Oregon. Oregon Department of Fish and Wildlife, Portland, OR, pp 5-52.

Beamesderfer RCP, Farr RA. 1997. Alternatives for the protection and restoration of sturgeons and their habitat. *Environ Biol Fish* 48:407-417.

Bendell BE, McNicol DK. 1995. The diet of insectivorous ducklings and the acidification of small Ontario lakes. *Can J Zool* 73:2044-2051.

Berlin WH, Hesselberg RJ, Mac MJ. 1981. Growth and mortality of fry of Lake Michigan lake trout during chronic exposure to PCBs and DDE. In: Chlorinated hydrocarbons as a factor in the reproduction and survival of lake trout (*Salvelinus namaycush*) in Lake Michigan. Technical Paper 105. US Fish and Wildlife Service, Washington, DC.

Beyer WN, Connor EE, Gerould S. 1994. Estimates of soil ingestion by wildlife. *J Wildl Manage* 58:375-382.

Biernacki M, Lovett-Doust J, Lovett-Doust L. 1997. Laboratory assay of sediment phytotoxicity using the macrophyte *Vallisneria spiralis*. *Environ Toxicol Chem* 16(3):472-478.

Birge WJ, Westerman AG, Spromberg JA. 2000. Comparative toxicology and risk assessment of amphibians. In: Sparling DW, Linder G, Bishop C, eds, *Ecotoxicology of amphibians and reptiles*. Society of Environ Toxicol Chem (SETAC), Pensacola, FL, pp 727-791.

Bjornn TC, Reiser DW. 1991. Habitat requirements of salmonids in streams. In: Meehan WR, ed, *Influences of forest and rangeland management on salmonid fishes and their habitats*. Vol 19. American Fisheries Society Special Publication, Bethesda, MD, pp 83-138.

Black JA, Birge WJ. 1980. An avoidance response bioassay for aquatic pollutants. Research rept. no. 123. Water Resources Research Unit, University of Kentucky, Lexington, KY.

Bleavins MR, Aulerich RJ. 1981. Feed consumption and food passage time in mink (*Mustela vison*) and European ferrets (*Mustela putorius furo*). *Lab Anim Sci* 31(3):268-269.

Bleavins MR, Aulerich RJ, Ringer RK. 1980. Polychlorinated biphenyls (Aroclors 1016 and 1242): Effects on survival and reproduction in mink and ferrets. *Arch Environ Contam Toxicol* 9:627-635.

Blue Ocean Tackle. 2011. Lead fishing weights: sinkers and lead heads [online]. Blue Ocean Tackle Inc., National City, CA. [Cited 5/16/11.] Available from: http://www.blueoceantackle.com/lead_fishing_weights.htm.

Boersma PD, Reichard SH, Van Buren AN, eds. 2006. Invasive species in the Pacific Northwest. University of Washington Press, Seattle, WA.

Bouton SN, Frederick PC, Spalding MG, McGill HC. 1999. Effects of chronic, low concentrations of dietary methylmercury on the behavior of juvenile great egrets. *Environ Toxicol Chem* 18(9):1934-1939.

Britton WM, Huston TM. 1973. Influence of polychlorinated biphenyls in the laying hen. *Poult Sci* 52:1620-1624.

Brooks KM. 1998. Literature review, computer model and assessment of the potential environmental risks associated with pentachlorophenol treated wood products used in aquatic environments. Prepared for Western Wood Preservers Institute, Vancouver, WA. Aquatic Environmental Services, Port Townsend, WA.

Brooks KM. 2004. The affects of dissolved copper on salmon and the environmental affects associated with the use of wood preservatives in aquatic environments. Prepared for Western Wood Preservers Institute, Vancouver, WA. Aquatic Environmental Services, Port Townsend, WA.

Brooks RP, Davis WJ. 1987. Habitat selection by breeding belted kingfishers. *Am Midl Nat* 117:63-70.

Broyles RH, Noveck MI. 1979. Uptake and distribution of 2,3,5,2',4',5'-hexachlorobiphenyl in fry of lake trout and chinook salmon and its effects on viability. *Toxicol Appl Pharmacol* 50:299-308.

Brunström B, Lund BE, Bergman A, Asplund L, Athanassiadis I, Athanasiadou M, Jensen S, Örborg J. 2001. Reproductive toxicity in mink (*Mustela vison*) chronically exposed to environmentally relevant polychlorinated biphenyl concentrations. *Environ Toxicol Chem* 20(10):2318-2327.

Brusca RC, Brusca GJ. 2003. Invertebrates. 2nd ed. Sinauer Associates, Sunderland, MA.

Buck J, Kaiser JL. 2011. Contaminant concentrations in osprey (*Pandion haliaetus*) eggs for Portland Harbor and surrounding areas: Data summary report. Prepared for the Portland Harbor Natural Resource Trustee Council. March 7, 2011. US Fish and Wildlife Service, Portland, OR; Kaiser and Associates Environmental Consulting, LLC, Seattle, WA.

Burkhard LP. 2009. Estimation of biota sediment accumulation factor (BSAF) from paired observations of chemical concentrations in biota and sediment. EPA/600/R-06/045. Ecological Risk Assessment Support Center, US Environmental Protection Agency, Cincinnati, OH.

Butler RW, Shepherd PCF, Lemon MJF. 2002. Site fidelity and local movements of migrating western sandpipers on the Fraser River estuary. *Wilson Bull* 114(4):485-490.

California DTSC. 2003. HERD ecological risk assessment Note 3: Calculation of an action level/preliminary cleanup goal for dibutyltin (DBT) in surface, ground, and sediment interstitial water for protection of saltwater aquatic life. 9/2/03. Human and Ecological Risk Division, Department of Toxic Substances Control, California Environmental Protection Agency, Sacramento, CA.

Carr RS, Long ER, Windom HL, Chapman DC, Thursby G, Sloane GM, Wolfe DA. 1996. Sediment quality assessment studies of Tampa Bay, Florida. *Environ Toxicol Chem* 15(7):1218-1231.

Carriere D, Fischer K, Peakall D, Angehrn P. 1986. Effects of dietary aluminum in combination with reduced calcium and phosphorous on the ring dove (*Streptopelia risoria*). *Wat Air Soil Pollut* 30:757-764.

CBFWA. 1996. Contamination ecology of selected fish and wildlife of the lower Columbia River. Report to the Bi-State Water Quality Program. Columbia Basin Fish & Wildlife Authority, Portland, OR.

Chapman PM, Fairbrother A, Brown D. 1998. A critical evaluation of safety (uncertainty) factors for ecological risk assessment. *Environ Toxicol Chem* 17(1):99-108.

City of Portland. 2008. Willamette River natural resources inventory: riparian corridors and wildlife habitat, Portland, Oregon. Proposed draft report, July 2008. City of Portland Bureau of Planning, Portland, OR.

Clemson University. 2011. Aquatic plant profiles: Soft rush (*Juncus effusus* var. *effusus*, *Juncus effusus* var. *solutus*) [online]. Cooperative Extension, Clemson University, Clemson, SC. [Cited 2/7/11.] Available from: http://www.clemson.edu/extension/horticulture/nursery/constructed_wetlands/plant_material/soft_rush.html.

Cleveland L, Buckler DR, Brumbaugh WG. 1991. Residue dynamics and effects of aluminum on growth and mortality in brook trout. *Environ Toxicol Chem* 10:243-248.

Cooke AS. 1972. The effects of DDT, dieldrin and 2,4-D on amphibian spawn and tadpoles. *Environ Pollut* 3:51-68.

Cooke SS. 1997. A field guide to the common wetland plants of Western Washington and Northwest Oregon. Seattle Audubon Society, Seattle, WA.

- Cooke SS, Azous AL. 1997. Characterization of Puget Sound basin palustrine wetland vegetation. Chapter 3. In: Azous AL, Horner RR, eds, Wetlands and urbanization: Implications for the future. Report of the Puget Sound wetlands and stormwater management research program. [Accessed 2007.] Available at: <<http://splash.metrokc.gov/wlr/basins/weturban.htm>>. King County Department of Natural Resources Seattle, WA.
<http://splash.metrokc.gov/wlr/basins/weturban.htm>.
- Corkran CC, Thoms C. 1996. Amphibians of Oregon, Washington and British Columbia. Lone Pine Publishing, Renton, WA.
- Cornwell GW. 1963. Observations on the breeding biology and behavior of a nesting population of belted kingfishers. *Condor* 65:426-431.
- Csuti B, O'Neil TA, Shaughnessy MM, Gaines EP, Hak JC. 2001. Atlas of Oregon wildlife: distribution, habitat, and natural history. 2nd ed. Oregon State University Press, Corvallis, OR.
- Cummins KW, Klug MJ. 1979. Feeding ecology of stream invertebrates. *Ann Rev Ecol Syst* 10:147-172.
- Dansereau M, Lariviere N, Du Tremblay D, Belanger D. 1999. Reproductive performance of two generations of female semidomesticated mink fed diets containing organic mercury contaminated freshwater fish. *Arch Environ Contam Toxicol* 36:221-226.
- Davison KL, Sell JL. 1974. DDT thins shells of eggs from mallard ducks maintained on *ad libitum* or controlled-feeding regimens. *Arch Environ Contam Toxicol* 2(3):222-232.
- de Smith M, Goodchild M, Longley P. 2008. Geospatial analysis: a comprehensive guide [online]. Matador, Leicester, UK. Available from: <http://www.spatialanalysisonline.com/output/>.
- DEA. 2003. Lower Willamette River multibeam bathymetric survey report. Prepared for Striplin Environmental Associates. David Evans and Associates, Inc., Portland, OR.
- Dean KE, Palachek RM, Noel JM, Warbritton R, Aufderheide J, Wireman J. 2004. Development of freshwater water-quality criteria for perchlorate. *Environ Toxicol Chem* 23(6):1441-1451.
- Dees LT. 1961. Sturgeons. Fishery leaflet #526. US Fish and Wildlife Service, Washington, DC.
- DeForest DK, Gensemer RW, Van Genderen EJ, Gorsuch JW. 2011a. Protectiveness of water quality criteria for copper in western United States waters relative to predicted olfactory responses in juvenile Pacific salmon. *Integr Environ Assess Manage* 7(3):336-347.
- DeForest DK, Meyer JS, Gensemer RW, Shephard BK, Adams WJ, Dwyer RL, Gorsuch JW, Van Genderen EJ. 2011b. Learned discourse: Are ambient water quality criteria for copper protective of olfactory impairment in fish? *Integr Environ Assess Manage* 7(1):145-146.

- DeVore JD, Grimes JT. 1993. Migration and distribution of white sturgeon in the Columbia River downstream from Bonneville Dam and in adjacent marine areas. In: Beamesderfer RC, Nigro AA, eds, Status and habitat requirements of the white sturgeon populations in the Columbia River downstream from McNary Dam. Final report of research. Vol I. Prepared for Bonneville Power Administration. Oregon Department of Fish and Wildlife, Portland, OR, pp 83-100.
- DeVore JD, James BW, Tracy CA, Hale DA. 1995. Dynamics and potential production of white sturgeon in the unpounded lower Columbia River. Trans Am Fish Soc 124:845-856.
- DeWitt JB. 1956. Chronic toxicity to quail and pheasants of some chlorinated insecticides. Agric Food Chem 4(10):863-866.
- Dickman M. 1968. The effect of grazing by tadpoles on the structure of a periphyton community. Ecology 49:1188-1190.
- Doe KG, Parker WR, Ponsford SJ, Vaughan JD. 1987. The acute and chronic toxicity of antimony to *Daphnia magna* and rainbow trout. Environmental Protection Service, Conservation and Protection, Environment Canada, Dartmouth, Nova Scotia.
- Domina DJ. 1997. Evaluation of the downstream migrant bypass system, T.W. Sullivan plant, Willamette Falls, Oregon: progress report for 1996. Portland General Electric Co., Portland, OR.
- Douben PET, ed. 2003. PAHs: an ecotoxicological perspective. John Wiley & Sons Ltd, Sharnbrook, Bedford, UK.
- Drilling N, Titman R, McKinney F. 2002. Mallard (*Anas platyrhynchos*). In: Poole A, ed, The birds of North America online. Cornell Laboratory of Ornithology, Ithaca, NY.
<http://bna.birds.cornell.edu/bna/species/658>.
- Dunning JB, ed. 1993. CRC handbook of avian body masses. CRC Press, Boca Raton, FL.
- Dunstone N, Birks JDS. 1987. The feeding ecology of mink (*Mustela vison*) in a coastal habitat. J Zool Lond 212:69-83.
- Dyer SD, White-Hull CE, Shephard BK. 2000. Assessments of chemical mixtures via toxicity reference values overpredict hazard to Ohio fish communities. Environ Sci Technol 34(12):2518-2524.
- Eagle TC, Whitman JS. 1987. Mink. In: Novak M, Baker JA, Obbard ME, Malloch B, eds, Wild furbearer management and conservation in North America. Ontario Ministry of Natural Resources, Toronto, ON, pp 615-624.
- Ecology. 2011a. Non-native invasive freshwater plants: Reed canarygrass (*Phalaris arundinacea*) [online]. Washington State Department of Ecology, Olympia, WA. [Cited 2/7/11.] Available from: <http://www.ecy.wa.gov/programs/wq/plants/weeds/aqua011.html>.

Ecology. 2011b. Shoreline plants: *Sagittaria cuneata* Sheld. and *Sagittaria latifolia* Willd. Duck potato, wapato, arrowhead [online]. Washington State Department of Ecology, Olympia, WA. [Cited 2/7/11.] Available from: <http://www.ecy.wa.gov/programs/wq/plants/plantid2/descriptions/sag.html>.

ECOTOX. 2009. ECOTOXicology database [online]. US Environmental Protection Agency. [Cited 2009.] Available from: http://www.epa.gov/ecotox/ecotox_home.htm.

Edwards EA, Krieger DA, Bacteller M, Maughan OE. 1982a. Habitat suitability index models: black crappie. FWS/OBS-82/10. US Fish and Wildlife Service, Washington, DC.

Edwards EA, Krieger DA, Gebhart G, Maughan OE. 1982b. Habitat suitability index models: white crappie. FWS/OBS-82/10.7. US Fish and Wildlife Service, Washington, DC.

Ehrlich PR, Dobkin DS, Wheye D. 1988. The birder's handbook. Simon and Schuster, New York, NY.

Elliott JE, Harris ML. 2001/2002. An ecotoxicological assessment of chlorinated hydrocarbon effects on bald eagle populations. Rev Toxicol 4:1-60.

Elliott JE, Henny CJ, Harris ML, Wilson LK, Norstrom RJ. 1999. Chlorinated hydrocarbons in livers of American mink (*Mustela vison*) and river otter (*Lutra canadensis*) from the Columbia and Fraser River basins, 1990-1992. Environ Monit Assess 57:229-252.

Ellis Ecological. 2000. A review and compilation of available information on fish use of the Portland Harbor, Portland, Oregon. Prepared for Port of Portland. Ellis Ecological Services, Inc., Estacada, OR.

Ellis Ecological. 2001. Juvenile salmonid residence time in Portland Harbor. Draft. Prepared for Lower Willamette Group. Ellis Ecological Services, Inc., Estacada, OR.

Emlen JM, Springman KR. 2007. Developing methods to assess and predict the population level effects of environmental contaminants. Integr Environ Assess Manage 3(2):157-165.

EPA. 1980a. Ambient water quality criteria for DDT. EPA 440/5-80-038. October 1980. US Environmental Protection Agency, Washington, DC.

EPA. 1980b. Ambient water quality criteria for naphthalene. EPA 440/5-80-059. October 1980. US Environmental Protection Agency, Washington, DC.

EPA. 1980c. Ambient water quality criteria for polychlorinated biphenyls. EPA 440/5-80-068. Office of Water, US Environmental Protection Agency, Washington, DC.

EPA. 1985. Ambient water quality criteria for cyanide - 1984. EPA 440/5-84-028. US Environmental Protection Agency, Washington, DC.

EPA. 1987. Ambient water quality criteria for zinc-1987. EPA-440/5-87-003. Office of Water, US Environmental Protection Agency, Washington, DC.

EPA. 1991. Ambient water quality criteria for tributyltin. EPA 822/R-91-100. March 1991. US Environmental Protection Agency, Duluth, MN.

EPA. 1992. Framework for ecological risk assessment. EPA/630/R-92/001. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 1993. Wildlife exposure factors handbook. EPA/600/R-93/187a. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

EPA. 1997. Ecological risk assessment guidance for Superfund: Process for designing and conducting ecological risk assessments. EPA/540/R-97/006. Interim final. Environmental Response Team, US Environmental Protection Agency, Edison, NJ.

EPA. 1998. Guidelines for ecological risk assessment. EPA/630/R-95/002 F. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 1999a. Issuance of final guidance: ecological risk assessment and risk management principles for Superfund sites. OSWER Directive 99285.7-28 P. October 7, 1999. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 1999b. USEPA contract laboratory program national functional guidelines for organic data review. EPA-540/R-99/008. October 1999. Office of Emergency and Remedial Response, US Environmental Protection Agency, Washington, DC.

EPA. 2000a. Guidance for assessing chemical contaminant data for use in fish advisories. Volume 1: Fish sampling and analysis. Third ed. EPA 823-B-00-007. November 2000. US Environmental Protection Agency, Washington, DC.

EPA. 2000b. Supplementary guidance for conducting health risk assessment of chemical mixtures. EPA/630/R-00/002. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 2001. Administrative Order on Consent for Remedial Investigation/Feasibility Study for Portland Harbor Superfund Site. US Environmental Protection Agency, Region 10, Portland, OR.

EPA. 2002a. Calculating upper confidence limits for exposure point concentrations at hazardous waste sites. Office of Emergency and Remedial Response, US Environmental Protection Agency, Washington, DC.

EPA. 2002b. Principles for managing contaminated sediment risks at hazardous waste sites. OSWER Directive 9285.6-08. Memorandum from M. Horinko dated February 12, 2002. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2003a. Ambient aquatic life water quality criteria for tributyltin (TBT) - final. EPA 822-R-03-031. US Environmental Protection Agency, Washington, DC.

EPA. 2003b. Analyses of laboratory and field studies of reproductive toxicity in birds exposed to dioxin-like compounds for use in ecological risk assessment. EPA/600/R-03/114F. National Center for Environmental Assessment, US Environmental Protection Agency, Cincinnati, OH.

EPA. 2003c. Procedures for the derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms: PAH mixtures. EPA-600-R-02-013. November 2003. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

EPA. 2004a. Considerations for developing problem formulations for ecological risk assessments conducted at contaminated sites under CERCLA. Environmental Response Team, US Environmental Protection Agency, Edison, NJ.

EPA. 2004b. An examination of EPA risk assessment principles and practices. EPA/100/B-04/001. US Environmental Protection Agency, Washington, DC.

EPA. 2004c. The incidence and severity of sediment contamination in surface waters of the United States. National Sediment Quality Survey, second ed. EPA 823-R-04-007. November 2004. US Environmental Protection Agency, Washington, DC.

EPA. 2005a. Contaminated sediment remediation guidance for hazardous waste sites. OSWER 9355.0-85. EPA-540-R-05-012. US Environmental Protection Agency, Office of Solid Waste and Emergency Response Washington, DC.

EPA. 2005b. Ecological soil screening levels for antimony. Interim final. OSWER Directive 9285.7-61. February 2005. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005c. Ecological soil screening levels for arsenic. Interim final. OSWER Directive 9285.7- 62. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005d. Ecological soil screening levels for cadmium. Interim final. OSWER Directive 9285.7-65. March 2005. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005e. Ecological soil screening levels for chromium. Interim final. OSWER Directive 9285.7-66. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005f. Ecological soil screening levels for lead. Interim final. OSWER Directive 9285.7-70. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2005g. EPA comments on technical memorandum: Provisional toxicity reference value selection for the Portland Harbor Preliminary Risk Assessment, March 18, 2005. Oregon Operations Office, US Environmental Protection Agency Region 10, Portland, OR.

EPA. 2005h. Predicting toxicity to amphipods from sediment chemistry. EPA/600/R-04/030. National Center for Environmental Assessment, US Environmental Protection Agency, Washington, DC.

EPA. 2006a. EPA comments on ecological risk assessment interpretive report: estimating risks to benthic organisms using predictive models based on sediment toxicity tests, and accompanying cover letter to Lower Willamette Group from C. Humphrey and E. Blischke dated July 6, 2006. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2006b. EPA letter dated October 31, 2006 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt) regarding Portland Harbor RI/FS Field Sampling Plan: Round 3 lamprey ammocoete toxicity testing field sampling plan. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2006c. Memorandum dated September 13, 2006 from D. Mount to ERAF tri-chairs (B. Pluta, M. Sprenger, V. Madden): Error in prior calculation of GLI Tier II SCV for m-xylene. National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, Duluth, MN.

EPA. 2006d. National recommended water quality criteria (4304T). November 2002. US Environmental Protection Agency, Washington, DC.

EPA. 2007a. Ecological soil screening levels for copper. Interim final. Revised February 2007. OSWER Directive 9285.7-68. Issued July 2006; revised February 2007. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2007b. Ecological soil screening levels for DDT and metabolites. Interim final. OSWER Directive 9285.7-57. April 2007. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2007c. Ecological soil screening levels for selenium. Interim final. OSWER Directive 9285.7-72. July 2007. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2007d. Ecological soil screening levels for zinc. Interim final. OSWER Directive 9285.7-73. June 2007. Office of Solid Waste and Emergency Response, US Environmental Protection Agency, Washington, DC.

EPA. 2007e. Framework for metals risk assessment. EPA 120/R-07/001. March 2007. Office of the Science Advisor, Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 2007f. ProUCL Version 4.0. Statistical software for environmental applications for data sets with and without nondetect observations [online]. Technical Support Center for Monitoring and Site Characterization, US Environmental Protection Agency. Updated 6/14/07. Available from: <http://www.epa.gov/nerlesd1/tsc/software.htm>.

EPA. 2008a. Calculation of aquatic biota toxicity reference values (TRVs) for petroleum alkanes, alkenes, cycloalkenes, BTEX and PAH compounds. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008b. EPA e-mail dated December 22, 2008 (Eric Blischke to Bob Wyatt, Rick Applegate, Jim McKenna) regarding fish TRVs for EPA submittal with attachment: EPA response to fish tissue-residue toxicity reference value reconciliation tables. Remedial Project Manager, US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008c. EPA e-mail dated June 27, 2008 (Burt Shephard to John Toll, Windward Environmental) regarding change of default ACR for use in water column TRV development. Risk Evaluation Unit, US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008d. EPA e-mail dated November 3, 2008 (Burt Shephard to Helle Andersen, Windward Environmental) regarding nutritional deficiency levels of zinc and copper in aquatic invertebrates. Risk Evaluation Unit, US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008e. EPA e-mail dated October 10, 2008 to Lower Willamette Group (from Eric Blischke to John Toll) regarding Portland Harbor RI/FS toxicity reference values -- elimination of aluminum from screening. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008f. EPA letter and attachment dated April 11, 2008 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt) regarding Portland Harbor RI/FS: Toxicity reference values for the baseline ecological risk assessment. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008g. EPA letter dated August 5, 2008 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt) regarding Portland Harbor RI/FS tissue TRV methodology, with attachments titled "Aquatic Tissue TRV response." Aquatic Biota Tissue TRV Derivation, and LWG Tissue TRV Response. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008h. EPA letter dated June 13, 2008 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt) regarding Portland Harbor RI/FS toxicity reference values methodology - aquatic biota tissue, with attachment titled "Aquatic Biota Tissue TRV Derivation". US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2008i. Framework for application of the toxicity equivalence methodology for polychlorinated dioxins, furans, and biphenyls in ecological risk assessment. EPA/100/R-08-004. June 2008. Risk Assessment Forum, US Environmental Protection Agency, Washington, DC.

EPA. 2008j. Problem formulation for the Baseline Ecological Risk Assessment at the Portland Harbor Site. Report and letter dated February 15, 2008 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt). US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2009a. EPA e-mail dated July 17, 2009, to Lower Willamette Group (from E. Blischke to R. Wyatt) regarding Portland Harbor RI/FS: interpretation of sediment bioassay results. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2009b. EPA e-mail dated July 31, 2009, to Lower Willamette Group (from E. Blischke to J. McKenna) regarding Portland Harbor RI/FS: benthic interpretation. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2009c. EPA e-mail dated March 17, 2009 (Eric Blischke to Bob Wyatt, Lower Willamette Group) regarding RI/FS agreement summary. Remedial Project Manager, US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2009d. EPA letter dated January 23, 2009 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna and R. Wyatt) regarding Portland Harbor RI/FS: Fish tissue-residue toxicity reference values for the baseline ecological risk assessment, with attachment titled "EPA evaluation of fish tissue-residue TRVs - behavioral endpoints". US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2009e. The national study of chemical residues in lake fish tissue. EPA-823-R-09-006. September 2009. Office of Water, US Environmental Protection Agency, Washington, DC.

EPA. 2010a. EPA letter and attachment dated April 21, 2010 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna) regarding Portland Harbor Superfund site: EPA direction to LWG on preliminary remediation goals (PRGs) for use in the Portland Harbor feasibility study. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA. 2010b. EPA letter and attachment dated September 27, 2010 to Lower Willamette Group (from E. Blischke and C. Humphrey to R. Wyatt) regarding Portland Harbor Superfund site: EPA comments on benthic risk evaluation. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.

EPA, USACE. 1998. Evaluation of dredged material proposed for discharge in waters of the U.S.- testing manual: Inland Testing Manual. EPA-823-B-98-004. US Environmental Protection Agency and US Army Corps of Engineers, Washington, DC.

Erickson RJ, Highland TL, Hockett JR, Leonard EN, Mattson VR, Mount DR. 2003. Effects of dietary copper, zinc, lead, cadmium, and arsenic on growth and survival of juvenile fish using live food organisms. Platform presentation at SETAC 24th annual meeting, Austin TX, 9-13 November 2003. Manuscript in prep.

Fagerlund F, Niemi A. 2003. Multi-constituent modelling of a gasoline spill using the T2VOC numerical simulator. Proceedings, TOUGH Symposium 2003, Lawrence Berkeley National Laboratory, Berkeley, CA. May 12-14, 2003.

Fairey R, Long ER, Roberts CA, Anderson BS, Phillips BM, Hunt JW, Puckett HM, Wilson CJ. 2001. An evaluation of methods for calculating mean sediment quality guideline quotients as indicators of contamination and acute toxicity to amphipods by chemical mixtures. *Environ Toxicol Chem* 20(10):276-2286.

Farg AM, Woodward DF, Goldstein JN, Brumbaugh W, Meyer JS. 1998. Concentrations of metals associated with mining waste in sediments biofilm, benthic macroinvertebrates, and fish from the Coeur d'Alene River Basin, Idaho. *Arch Environ Contam Toxicol* 34:119-127.

Farr RA, Ward DL. 1992. Fishes of the lower Willamette River near Portland, Oregon. Oregon Department of Fish and Wildlife, Clackamas, OR.

Farr RA, Ward DL. 1993. Fishes of the lower Willamette River near Portland, Oregon. *Northwest Sci* 67(1):16-22.

Field J, Norton S, MacDonald D, Severn C, Ingersoll C. 1999a. Beyond thresholds: using logistic regression models to estimate the probability of toxicity from sediment chemistry. SETAC poster.

Field LJ, MacDonald DD, Norton SB, Severn CG, Ingersoll CG. 1999b. Evaluating sediment chemistry and toxicity data using logistic regression modeling. *Environ Toxicol Chem* 18:1311-1322.

Field LJ, MacDonald DD, Norton SB, Ingersoll CG, Severn CG, Smorong D, Lindscoog R. 2002. Predicting amphipod toxicity from sediment chemistry using logistic regression models. *Environ Toxicol Chem* 21(9):1993–2005.

Fisher JP, Spitsbergen JM, Bush B, Jahan-Parwar B. 1994. Effect of embryonic PCB exposure on hatching success, survival, growth and developmental behavior in landlocked Atlantic salmon, *Salmo salar*. In: Gorsuch JW, Dwyer FJ, Ingersoll CG, La Point TW, eds, *Environmental toxicology and risk assessment*. Vol 2. ASTM STP 1216. American Society for Testing and Materials, Philadelphia, PA, pp 298-314.

Fishman. 1999. Eastbank riverfront (phase I) floating walkway fish predation study: data summary, spring 1999 sampling season. Prepared for Portland Development Commission. Fishman Environmental Services, Portland, OR.

Fletcher JS. 1990. Use of algae versus vascular plants to test for chemical toxicity. In: Wang W, Gorsuch JW, Lower WR, eds, *Plants for toxicity assessment*. ASTM STP 1091. American Society for Testing and Materials, Philadelphia, PA, pp 33-39.

Folmar LC. 1976. Overt avoidance reaction of rainbow trout fry to nine herbicides. *Bull Environ Contam Toxicol* 15:509.

Forbes TL, Forbes VE, Giessing A, Hansen R, Kure LK. 1998. Relative role of pore water versus ingested sediment in bioavailability of organic contaminants in marine sediments. *Environ Toxicol Chem* 17(12):2453-2462.

Forbes VE, Calow P. 2002. Population growth rate as a basis for ecological risk assessment of toxic chemicals. *Phil Trans R Soc Lond B* 357:1299-1306.

Foster CA, Boatner JR. 2002. 2000 Lower Willamette and Clackamas River steelhead and Willamette River white sturgeon and shad recreational fisheries. In: 2000 Willamette River spring chinook salmon run, fisheries, and passage at Willamette Falls. Oregon Department of Fish and Wildlife, Clackamas, OR.

Friesen TA, Ward DL. 1996. Status and condition of fish assemblages in streams of the Tualatin River Basin, Oregon. *Northwest Sci* 70:120-31.

Friesen TA, ed. 2005. Biology, behavior, and resources of resident and anadromous fish in the Lower Willamette River. Final report of research, 2000-2004. Prepared for City of Portland. Oregon Department of Fish and Wildlife, Clackamas, OR.

Fuchsman PC, Barber TR, Bock MJ. 2007. Effectiveness of various exposure metrics in defining dose-response relationships for mink (*Mustela vison*) exposed to polychlorinated biphenyls. *Arch Environ Contam Toxicol* 54:130-144.

Garrett MG, Watson JW, Anthony RG. 1993. Bald eagle home range and habitat use in the Columbia River estuary. *J Wildl Manage* 57(1):19-27.

- Gatlin DM, III, Wilson RP. 1986. Dietary copper requirement of fingerling channel catfish. *Aquaculture* 54:277-285.
- George EL, Hadley WF. 1979. Food and habitat partitioning between rock bass (*Ambloplites rupestris*) and smallmouth bass (*Micropterus dolomieu*) young of the year. *Trans Am Fish Soc* 108:253-261.
- Giattina JD, Garton RR, Stevens DG. 1982. Avoidance of copper and nickel by rainbow trout as monitored by a computer-based data acquisition system. *Trans Am Fish Soc* 111:491-504.
- Golub MS, Gershwin ME, Donald JM, Negri S, Keen CL. 1987. Maternal and developmental toxicity of chronic aluminum exposure in mice. *Fund Appl Toxicol* 8(5):346-357.
- Guard BJ. 1995. Wetland plants of Oregon and Washington. Lone Pine Publishing, Renton, WA.
- Hansen JA, Woodward DF, Little EE, DeLonay AJ, Bergman HL. 1999a. Behavioral avoidance: possible mechanism for explaining abundance and distribution of trout species in a metal-impacted river. *Environ Toxicol Chem* 18(2):313-317.
- Hansen JA, Rose JD, Jenkins RA, Gerow KG, Bergman HL. 1999b. Chinook salmon (*Oncorhynchus tshawytscha*) and rainbow trout (*Oncorhynchus mykiss*) exposed to copper: neurophysiological and histological effects on the olfactory system. *Environ Toxicol Chem* 18(9):1979-1991.
- Hansen JA, Marr JCA, Lipton J, Cacela D, Bergman HL. 1999c. Differences in neurobehavioral responses of chinook salmon (*Oncorhynchus tshawytscha*) and rainbow trout (*Oncorhynchus mykiss*) exposed to copper and cobalt: behavioral avoidance. *Environ Toxicol Chem* 18(9):1972-1978.
- Hara TJ, Law YMC, MacDonald S. 1976. Effects of mercury and copper on the olfactory response in rainbow trout, (*Salmo gairdneri*). *J Fish Res Bd Can* 33:1568-1573.
- Hare L, Tessier A, Warren L. 2001. Cadmium accumulation by invertebrates living at the sediment-water interface. *Environ Toxicol Chem* 20(4):880-889.
- Hayes MP, Tennant MR. 1985. Diet and feeding behavior of the California red-legged frog *Rana aurora draytonii* (Ranidae). *Southw Naturalist* 30(4):601-605.
- Healey MC. 1991. Life history of chinook salmon (*Oncorhynchus tshawytscha*). In: Groot C, Margolis L, eds, Pacific salmon life histories. UBC Press, Vancouver, BC, pp 311-394.
- Heath RG, Spann JW, Kreitzer JF. 1969. Marked DDE impairment of mallard reproduction in controlled studies. *Nature* 224:47-48.

Heaton SN, Bursian SJ, Giesy JP, Tillitt DE, Render JA, Jones PD, Verbrugge DA, Kubiak TJ, Aulerich RJ. 1995. Dietary exposure of mink to carp from Saginaw Bay, Michigan. 1. Effects on reproduction and survival, and the potential risks to wild mink populations. *Arch Environ Contam Toxicol* 28:334-343.

Heinz GH. 1974. Effects of low dietary levels of methyl mercury on mallard reproduction. *Bull Environ Contam Toxicol* 11(4):386-392.

Heinz GH. 1975. Effects of methylmercury on approach and avoidance behavior of mallard ducklings. *Bull Environ Contam Toxicol* 13(5):554-564.

Heinz GH. 1976. Methylmercury: second-year feeding effects on mallard reproduction and duckling behavior. *J Wildl Manage* 40(1):82-90.

Heinz GH. 1979. Methylmercury: reproductive and behavioral effects on three generations of mallard ducks. *J Wildl Manage* 43(2):394-401.

Henny CJ, Grove RA, Hedstrom OR. 1996. A field evaluation of mink and river otter on the lower Columbia River and the influence of environmental contaminants. Final report. Prepared for Lower Columbia River Bi-State Water Quality Program. Northwest Research Station, Forest and Rangeland Ecosystem Science Center, National Biological Service, Corvallis, OR.

Henny CJ, Kaiser JL, Grove RA, Bentley VR, Elliott JE. 2003. Biomagnification factors (fish to osprey eggs from Willamette River, Oregon, USA) for PCDDs, PCDFs, PCBs and OC pesticides. *Environ Monit Assess* 84:275-315.

Henny CJ, Grove RA, Kaiser JL, Bentley VR. 2004. An evaluation of osprey eggs to determine spatial residue patterns and effects of contaminants along the lower Columbia River, USA. In: Chancellor R, Meyburg B, eds, *Raptors worldwide*. WWGBP/MME, Budapest, Hungary, pp 369-388.

Henny CJ, Kaiser JL, Grove RA. 2009. PCDDs, PCDFs, PCBs, OC pesticides and mercury in fish and osprey eggs from Willamette River, Oregon (1993, 2001 and 2006) with calculated biomagnification factors. *Ecotoxicology* 18:151-173.

Hilsenhoff WL. 1987. An improved biotic index of organic stream pollution. *Great Lakes Entomol* 20:31-40.

Hinck JE, Schmitt CJ, Bartish TM, Denslow ND, Blazer VS, Anderson PJ, Coyle JJ, Dethloff GM, Tillitt DE. 2004. Biomonitoring of environmental status and trends (BEST) program: environmental contaminants and their effects on fish in the Columbia River basin. Report no. 2004-5154. Columbia Environmental Research Center, US Geological Survey, Columbia, MO.

Hitchcock CL, Cronquist A. 1973. *Flora of the Pacific Northwest*. University of Washington Press, Seattle, WA.

- Hoffman B. 2005. Effect of advective pore water flow on degradation of organic matter in permeable sandy sediment: a study of fresh and brackish water. Masters thesis. Linkopings University, Norrkoping, Sweden. 2005.
- Hoffman DJ, Rattner BA, Burton Jr. GA, Cairns Jr. J. 1995. Handbook of ecotoxicology. Lewis Publishers, Boca Raton, FL.
- Holcombe GW, Benoit DA, Leonard EN, McKim JM. 1976. Long-term effects of lead exposure on three generations of brook trout (*Salvelinus fontinalis*). J Fish Res Bd Can 33:1731-1741.
- Hope BK. 2003. A basin-specific aquatic food web biomagnification model for estimation of mercury target levels. Environ Toxicol Chem 22(10):2525-2537.
- Hopkins CL, Solly SRB, Ritchie AR. 1969. DDT in trout and its possible effect on reproductive potential. NZ J Mar Freshw Res 3:220-229.
- Horness BH, Lomax DP, Johnson LL, Myers MS, Pierce SM, Collier TK. 1998. Sediment quality thresholds: estimates from hockey stick regression of liver lesion prevalence in English sole (*Pleuronectes vetulus*). Environ Toxicol Chem 17:872-882.
- Hornshaw TC, Aulerich RJ, Johnson HE. 1983. Feeding Great Lakes fish to mink: effects on mink and accumulation and elimination of PCBs by mink. J Toxicol Environ Health 11:933-946.
- Hornshaw TC, Safronoff J, Ringer RK, Aulerich RJ. 1986. LC50 test results in polychlorinated biphenyl-fed mink: age, season, and diet comparisons. Arch Environ Contam Toxicol 15:717-723.
- Hough JL, Baird MB, Sfeir GT, Pacini CS, Darrow D, Wheelock C. 1993. Benzo(a)pyrene enhances atherosclerosis in white carneau and show racer pigeons. Arterioscler Thromb 13:1721-1727.
- Houston CS, Bowen DE, Jr. 2001. Upland sandpiper (*Bartramia longicauda*). In: Poole A, ed, The birds of North America online. Cornell Laboratory of Ornithology, Ithaca, NY. <http://bna.birds.cornell.edu/bna/species/580doi:10.2173/bna.580>.
- Hubert TD. 2003. Environmental fate and effects of the lampricide TFM: a review. J Great Lakes Res 29(Suppl 1):456-474.
- Hudson RA, Austerberry CF, Bagshaw JC. 1981. Phthalate ester hydrolases and phthalate ester toxicity in synchronously developing larvae of the brine shrimp (*Artemia*). Life Sci 29:1865-1872.
- Hudson RH, Tucker RK, Haegele MA. 1984. Handbook of toxicity of pesticides to wildlife. Resource publication 153. 2nd ed. US Fish and Wildlife Service, Washington, DC.
- Huettel M, Rusch A. 2000. Transport and degradation of phytoplankton in permeable sediments. Limnol Oceanogr 45(3):534-549.

Hughes RM, Gammon JR. 1987. Longitudinal changes in fish assemblages and water quality in the Willamette River, Oregon. *Trans Am Fish Soc* 116:196-209.

Hugla JL, Thome JP. 1999. Effects of polychlorinated biphenyls on liver ultrastructure, hepatic monooxygenases, and reproductive success in the barbel. *Ecotox Environ Saf* 42:265-273.

Humphrey C. 2011. Personal communication on February 25, 2011 (re: Portland Harbor AOC for RI/FS Schedule). US Environmental Protection Agency, Seattle, WA.

Integral. 2004a. Portland Harbor RI/FS round 2a field sampling plan: surface water sampling. Prepared for Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA.

Integral. 2004b. Portland Harbor RI/FS: Round 1 site characterization summary report. Draft. Prepared for Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA.

Integral. 2006a. Portland Harbor RI/FS: Round 2 groundwater pathway assessment, transition zone water site characterization summary report. IC-06-0020. Prepared for the Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA.

Integral. 2006b. Portland Harbor RI/FS: Round 3a field sampling plan, surface water sampling. Draft. IC-06-0012. Prepared for the Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA.

Integral, Windward, Kennedy/Jenks, Anchor, Groundwater Solutions. 2004a. Portland Harbor RI/FS programmatic work plan. Appendix B: ecological risk assessment. Prepared for Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA; Windward Environmental LLC, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR; Anchor Environmental, LLC, Seattle, WA; Groundwater Solutions, Inc., Portland, OR.

Integral, Windward, Kennedy/Jenks, Anchor, Groundwater Solutions. 2004b. Portland Harbor RI/FS programmatic work plan. Prepared for Lower Willamette Group. April 23, 2004. Integral Consulting, Inc., Mercer Island, WA; Windward Environmental LLC, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR; Anchor Environmental, LLC, Seattle, WA; Groundwater Solutions, Inc., Portland, OR.

Integral, Windward, Kennedy/Jenks, Anchor. 2007. Portland Harbor RI/FS: Comprehensive round 2 site characterization summary and data gaps analysis report, plus addenda. IC07-0004. Prepared for Lower Willamette Group. Integral Consulting, Inc., Mercer Island, WA; Windward Environmental LLC, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR; Anchor Environmental, LLC, Portland, OR.

Integral, Windward, Kennedy/Jenks, Anchor QEA. 2011. Portland Harbor RI/FS: Remedial investigation report. IC09-0003. Draft final. Prepared for Lower Willamette Group. August 2011. Integral Consulting, Inc., Mercer Island, WA; Windward Environmental LLC, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR; Anchor QEA, LLC, Portland, OR.

Isaacs FB, Anthony RG. 2001. Bald eagle nest locations and history of use in Oregon and the Washington portion of the Columbia River recovery zone, 1972 through 2001. Oregon Cooperative Fish and Wildlife Research Unit, Oregon State University, Corvallis, OR.

Isensee AR. 1978. Bioaccumulation of 2,3,7,8-tetrachlorodibenzi-*para*-dioxin. Ecol Bull (Stockholm) 27:255-262.

Ishida M, Suyama K, Adachi S, Hoshino T. 1982. Distribution of orally administered di-ethylhexyl phthalate in laying hens. Poult Sci 61:262-267.

Jahn LA, Anderson RV. 1986. The ecology of pools 19 and 20, upper Mississippi River : a community profile. National Wetlands Research Center, Research and Development, Fish and Wildlife Service, US Department of the Interior, Washington, DC.

Jehl JR, Jr. 1973. Breeding biology and systematic relationships of the stilt sandpiper. Wilson Bull 85:114-147.

Jensen S, Kihlstrom JE, Olsson M, Lundberg C, Orberg J. 1977. Effects of PCB and DDT on mink (*Mustela vison*) during the reproductive season. Ambio 6(4):239.

Johnson L, Ylitalo GM, Myers MS, Anulacion BF, Buzitis J, Reichert WL, Collier TK. 2009. Polycyclic aromatic hydrocarbons and fish health indicators in the marine ecosystem in Kitimat, British Columbia. NOAA Tech Memo NMFS-NWFSC-98. March 2009. Northwest Fisheries Science Center, National Marine Fisheries Service, Seattle, WA.

Johnson LL. 2000. An analysis in support of sediment quality thresholds for polycyclic aromatic hydrocarbons (PAHs) to protect estuarine fish. Memorandum to Rachel Friedman, Steven Landino, dated July 24, 2000. Environmental Conservation Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA.

Johnson LL, Collier TK, Stein JE. 2002. An analysis in support of sediment quality thresholds for polycyclic aromatic hydrocarbons (PAHs) to protect estuarine fish. Aquat Conserv Mar Freshw Ecosys 12(5):517-538.

Johnson LL, Landahl JT. 1994. Chemical contaminants, liver disease, and mortality rates in English sole (*Pleuronectes vetulus*). Ecol Applic 4:59-68.

Julliard AK, Saucier D, Astic L. 1996. Time-course of apoptosis in the olfactory epithelium of rainbow trout exposed to a low copper level. Tissue Cell 28:367-377.

Kaiser-Hill. 2006. RCRA facility investigation - remedial investigation/corrective measures study - feasibility study report for the Rocky Flats Environmental Technology site. Appendix A - Comprehensive risk assessment. Volume 1 of 15: Executive summary. Prepared for US Department of Energy. June 2006. Kaiser-Hill Company, L.L.C., Broomfield, CO.

- Kang J-C, Kim S-G, Jang S-W. 2005. Growth and hematological changes of rockfish, *Sebastes schlegeli* (Hilgendorf) exposed to dietary Cu and Cd. J World Aquacult Soc 36(2):188-195.
- Kennedy/Jenks. 2011. Portland Harbor RI, Appendix F: Baseline human health risk assessment. Draft final. Prepared for the Lower Willamette Group. May 2, 2011. Kennedy/Jenks Consultants, Portland, OR.
- Kihlstrom JE, Olsson M, Jensen S, Johansson A, Ahlbom J, Bergman A. 1992. Effects of PCB and different fractions of PCB on the reproduction of the mink (*Mustela vison*). Ambio 2(8):563-569.
- Kim S-G, Kim J-W, Kang J-C. 2004. Effect of dietary cadmium on growth and haematological parameters of juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). Aquacult Res 35:80-86.
- Kitchen DW, Hunt GS. 1969. Brood habitat of the hooded merganser. J Wildl Manage 33:605-609.
- Klima J, Jehl JR, Jr. 1998. Stilt sandpiper (*Calidris himantopus*). In: Poole A, ed, The birds of North America online. Cornell Laboratory of Ornithology, Ithaca, NY.
<http://bna.birds.cornell.edu/bna/species/341doi:10.2173/bna.341>.
- Kostow K. 2002a. Oregon lampreys: natural history status and analysis of management issues. February 25, 2002. Oregon Department of Fish and Wildlife, Portland, OR.
- Kostow K. 2002b. Oregon lampreys: natural history status and problem analysis. February 25, 2002. Oregon Department of Fish and Wildlife, Portland, OR.
- Krantzberg G. 1985. The influence of bioturbation on physical, chemical and biological parameters in aquatic environments. A review. Environ Pollut (Series A) 39(2):99-122.
- Krieger DA, Terrell JW, Nelson PC. 1983. Habitat suitability information: Yellow perch. FWS/OBS-83/10.55. US Fish and Wildlife Service, Washington, DC.
- Lall SP, Hines JA. 1987. Iron and copper requirements of Atlantic salmon (*Salmo salar*). In: International Symposium on Feeding and Nutrition in Fish, 23-27 August 1987, Bergen, Norway: Abstracts. p 48.
- Lancot RB, Laredo CD. 1994. Buff-breasted sandpiper (*Tryngites subruficollis*). In: Poole A, ed, The birds of North America online. Cornell Laboratory of Ornithology, Ithaca, NY.
<http://bna.birds.cornell.edu/bna/species/091/articles/introduction>.
- Landahl JT, McCain BB, Myers MS, Rhodes JD, Brown DW. 1990. Consistent associations between hepatic lesions in English sole (*Parophrys vetulus*) and polycyclic aromatic hydrocarbons in bottom sediment. Environ Health Perspect 89:195-203.
- Lannoo MJ. 2005. Amphibian declines: the conservation status of United States species. University of California Press, Berkeley, CA.

Lee H, II, Swartz RC. 1980. Chapter 29. Biological processes affecting the distribution of pollutants in marine sediments, Part II. biodeposition and bioturbation. In: Baker RA, ed, Contaminants and sediments. Vol 2. Ann Arbor Science Publishers, Ann Arbor, MI, pp 555-606.

Lewis MA, Weber DE, Stanley RS, Moore JC. 2001. The relevance of rooted vascular plants as indicators of estuarine sediment quality. *Arch Environ Contam Toxicol* 40:25-34.

Lim C, Yildirim-Aksoy M, Klesius PH. 2008. Nutrition and disease resistance in fish. In: Cyrino JEP, Bureau DP, Kapoor BG, eds, Feeding and digestive functions of fishes. Science Publishers, Inc, Enfield, NH, pp 479-545.

Lincer JL. 1975. DDE-induced eggshell-thinning in the American kestrel: a comparison of the field situation and laboratory results. *J Appl Ecol* 12(3):781-793.

Long ER, Field LJ, MacDonald DD. 1998. Predicting toxicity in marine sediments with numerical sediment quality guidelines. *Environ Toxicol Chem* 17(4):714-727.

Long ER, Ingersoll CG, MacDonald DD. 2006. Calculation and uses of mean sediment quality guideline quotients: a critical review. *Environ Sci Technol* 40(6):1726-1736.

Lorentzen M, Maage A, Julshamn K. 1998. Supplementing copper to a fishmeal-based diet fed to Atlantic salmon parr affects liver copper and selenium concentrations. *Aquacult Nutr* 4:67-72.

Lotufo GR, Landrum PF, Gedeon ML, Tigue EA, Herche LR. 2000. Comparative toxicity and toxicokinetics of DDT and its major metabolites in freshwater amphipods. *Environ Toxicol Chem* 19(2):368-379.

Lundholm CE. 1997. DDE-induced eggshell thinning in birds: effects of p,p'-DDE on the calcium and prostaglandin metabolism of the eggshell gland. *Comp Biol Physiol* 118C(2):113-128.

Luoma SN, Rainbow PS. 2008. Metal contamination in aquatic environments: science and lateral management. Cambridge University Press, Cambridge, UK.

LWG. 2010. General responses to EPA's non-directed comment key issues on the draft Baseline Ecological Risk Assessment. November 18, 2010. Lower Willamette Group, Portland, OR.

LWG. 2011a. Letter and attachments dated January 12, 2011 from B. Wyatt to C. Humphrey, EPA Region 10: response to December 21, 2010 EPA letter on the status of the Portland Harbor Feasibility Study; September 27, 2010 EPA letter on the Benthic Risk Evaluation; and December 8, 2010 EPA letter on general responses to EPA non-directed RI, BHHRA and BERA comments. Lower Willamette Group, Portland, OR.

LWG. 2011b. Letter dated March 9, 2011, from Bob Wyatt to Chip Humphrey and Kristine Koch, EPA Region 10: response to EPA's February 25, 2011, response to LWG proposed project schedule presented on February 2, 2011. Draft. Lower Willamette Group, Portland, OR.

Mac MJ, Edsall CC, Seelye JG. 1985. Survival of lake trout eggs and fry reared in water from the upper Great Lakes. *J Great Lakes Res* 11(4):520-529.

MacDonald DD, Landrum PF. 2008. An evaluation of the approach for assessing risks to the benthic invertebrate community at the Portland Harbor Superfund site. Preliminary draft. Prepared for US Environmental Protection Agency. September 2008. MacDonald Environmental Sciences, Ltd., Nanaimo, BC, and Landrum and Associates, Ann Arbor, MI.

MacDonald DD, Ingersoll CG, Berger TA. 2000. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 39(5):20-31.

MacDonald Environmental. 2002. Calcasieu Estuary remedial investigation/feasibility study (RI/FS): Baseline ecological risk assessment (BERA). Prepared for CDM Federal Programs Corp. September 2002. MacDonald Environmental Services, Ltd., Nanaimo, BC.

MacKenzie KM, Angevine DM. 1981. Infertility in mice exposed *in utero* to benzo(a)pyrene. *Biol Reprod* 24:183-191.

Malins DC, Anderson KM, Stegeman JJ, Jaruga P, Green VM, Gilman NK, Dizdaroglu M. 2006. Biomarkers signal contaminant effects on the organs of English sole (*Parophrys vetulus*) from Puget Sound. *Environ Health Perspect* 14(6):823-829.

Marshall DB, Hunter MG, Contreras AL, eds. 2003. *Birds of Oregon*. Oregon State University Press, Corvallis, OR.

Massey JB. 1967. The downstream migration of juvenile anadromous fish at Willamette Falls, Oregon. June 1, 1965 through November 30, 1966. Fishery Division, Oregon State Game Commission, Portland, OR.

Matta MB, Linse J, Cairncross C, Francendese L, Kocan RM. 2001. Reproductive and transgenerational effects of methylmercury or Aroclor 1268 on *Fundulus heteroclitus*. *Environ Toxicol Chem* 20(2):327-335.

Maxson SJ, Oring LW. 1980. Breeding season time and energy budgets of the polyandrous spotted sandpiper. *Behaviour* 74:200-263.

McCabe GT, Hinton SA, Emmett RL, Sandford BP. 1997. Benthic invertebrates and sediment characteristics in main channel habitats in the lower Columbia River. *Northwest Sci* 71(1):45-55.

McFarland VA. 1995. Evaluation of field-generated accumulation factors for predicting the bioaccumulation potential of sediment-associated PAH compounds. Long-term effects of dredging operations program technical report D-95-2. Waterways Experiment Station, US Army Corps of Engineers, Vicksburg, MS.

McIntyre JK, Baldwin DH, Meador JP, Scholtz NL. 2008. Chemosensory deprivation in juvenile coho salmon exposed to dissolved copper under varying water chemistry conditions. *Environ Sci Technol* 42:1352-1358.

McMahon RF, Bogen AE. 2001. Mollusca: Bivalvia. In: Thorp JH, Covich AP, eds, *Ecology and classification of North American freshwater invertebrates*. Academic Press, San Diego, CA, pp 331-431.

McMahon TE, Gebhart G, Maughan OE, Nelson PC. 1984a. Habitat suitability index models and instream flow suitability curves: Warmouth. FWS/OBS-82/10.67 [online]. US Fish and Wildlife Service. [Cited 3/28/02.] Available from: <http://www.nwrc.gov/wdb/pub/hsi/hsi-067.pdf>.

McMahon TE, Terrell JW, Nelson PC. 1984b. Habitat suitability information: Walleye. FWS/OBS-82/10.56. US Fish and Wildlife Service, Washington, DC.

Meador JP, Collier TK, Stein JE. 2002a. Determination of a tissue and sediment threshold for tributyltin to protect prey species of juvenile salmonids listed under the US Endangered Species Act. *Aquat Conserv Mar Freshw Ecosys* 12:539-551.

Meador JP, Collier TK, Stein JE. 2002b. Use of tissue and sediment-based threshold concentrations of polychlorinated biphenyls (PCBs) to protect juvenile salmonids listed under the US Endangered Species Act. *Aquat Conserv Mar Freshw Ecosys* 12:493-516.

Meador JP, Sommers FC, Ylitalo GM, Sloan CA. 2006. Altered growth and related physiological responses in juvenile chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). *Can J Fish Aquat Sci* 63:2364-2376.

Mehrle PM, Mayer FL. 1976. Di-2-ethylhexyl phthalate: residue dynamics and biological effects in rainbow trout and fathead minnows. In: Hemphill DD, ed, *Proceedings of University of Missouri's 10th Annual Conference on Trace Substances in Environmental Health*, June 8-10, Columbia, MO, pp 519-524.

Melquist WE, Whitman JS, Hornocker MG. 1981. Resource partitioning and coexistence of sympatric mink and river otter populations. In: Chapman JA, Pursley D, eds, *Worldwide Furbearer Conference Proceedings*. Worldwide Furbearer Conference Inc, Frostburg, MD, pp 187-220.

Mendenhall VM, Klaas EE, McLane MAR. 1983. Breeding success of barn owls (*Tyto alba*) fed low levels of DDE and dieldrin. *Arch Environ Contam Toxicol* 12:235-240.

Merritt RW, Cummins KW, Burton TW. 1984. The role of aquatic insects in the processing and cycling of nutrients. In: Resh VH, Rosenberg DM, eds, *The ecology of aquatic insects*. Praeger Publishers, New York, NY, pp 134-163.

- Meyer JS, Adams WJ. 2010. Relationship between biotic ligand model-based water quality criteria and avoidance and olfactory responses to copper by fish. *Environ Toxicol Chem* 29(9):2096-2103.
- Meyer JS, Adams WJ, Brix KV, Luoma SN, Mount DR, Stubblefield WA, Wood CM, eds. 2005. Toxicity of dietborne metals to aquatic organisms. Chapter 1, introduction, pp. 1-11. Society of Environmental Toxicology and Chemistry (SETAC), Pensacola, FL.
- Mezin LC, Hale RC. 2000. Effects of contaminated sediment on the epidermis of mummichog, *Fundulus heteroclitus*. *Environ Toxicol Chem* 19(11):2779-2787.
- Moore JW, Mallatt JM. 1980. Feeding of larval lamprey. *Can J Fish Aquat Sci* 37:1658-1664.
- Murai T, Andrews JW, Smith RG, Jr. 1981. Effects of dietary copper on channel catfish. *Aquaculture* 22:353-357.
- Myers MS, Stehr CM, Olson OP, Johnson LL, McCain BB, Chan SL, Varanasi U. 1994. Relationships between toxicopathic hepatic lesions and exposure to chemical contaminants in English sole (*Pleuronectes vetulus*), starry flounder (*Platichthys stellatus*), and white croaker (*Genyonemus lineatus*) from selected marine sites on the Pacific coast, USA. *Environ Health Perspect* 102:200-215.
- Nagy KA. 1987. Field metabolic rate and food requirement scaling in mammals and birds. *Ecol Monogr* 57(2):111-128.
- Nagy KA. 2001. Food requirements of wild animals: predictive equations for free-living mammals, reptiles, and birds. *Nutrition Abstracts and Reviews Series B: Livestock Feeds and Feeding* 71(10):21R-31R.
- Nakayama K, Oshima Y, Nagafuchi K, Hano T, Shimasaki Y, Honjo T. 2005. Early-life-stage toxicity in offspring from exposed parent medaka, *Oryzias latipes*, to mixtures of tributyltin and polychlorinated biphenyls. *Environ Toxicol Chem* 24(3):591-596.
- Niimi AJ. 1983. Biological and toxicological effects of environmental contaminants in fish and their eggs. *Can J Fish Aquat Sci* 40:306-312.
- NMFS. 1996. Status review of west coast steelhead. NOAA Technical Memorandum NMFS-NWFSC-27. National Marine Fisheries Service, Seattle, WA.
- North JA, Burner LC, Cunningham BC, Farr RA, Friesen TA, Harrington JC, Takata HK, Ward DL. 2002. Relationships between bank treatment/nearshore development and anadromous/resident fish in the lower Willamette River. Annual progress report, May 2000-June 2001. Draft. Prepared for City of Portland. Oregon Department of Fish and Wildlife, Clackamas, OR.

Nosek JA, Craven SR, Sullivan JR, Hurley SS, Peterson RE. 1992. Toxicity and reproductive effects of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in ring-necked pheasant hens. *J Toxicol Environ Health* 35:187-198.

O'Neill SM, West JE, Hoeman JC. 1998. Spatial trends in the concentration of polychlorinated biphenyls (PCBs) in chinook (*Oncorhynchus tshawytscha*) and coho salmon (*O. kisutch*) in Puget Sound and factors affecting PCB accumulation: Results from the Puget Sound Ambient Monitoring Program. *Puget Sound Research* '98:312-328.

ODEQ. 1998. (Updated 2001.) Guidance for ecological risk assessment: levels I, II, III, IV. Waste Management and Cleanup Division, Oregon Department of Environmental Quality, Portland, OR.

ODEQ. 2005. Laboratory analytical storage and retrieval (LASAR) database. Data for Willamette River At Sp&S RR Bridge (Portland) [online]. DEQ Online, Oregon Department of Environmental Quality, Portland, OR. Updated 6/24/05. Available from: <http://www.deq.state.or.us/wq/lasar/LasarHome.htm>.

ODEQ. 2006. Water pollution, Division 41, water quality standards: beneficial uses, policies and criteria for Oregon. OAR 340-041-0001 [online]. Oregon Department of Environmental Quality, Salem, OR. Updated 12/15/06. [Cited 1/26/07.] Available from: <http://www.deq.state.or.us/regulations/rules.htm>.

ODFW. 2001. Fisheries management and evaluation plan: Upper Willamette River spring chinook in freshwater fisheries of the Willamette basin and lower Columbia River mainstem. February 2001. Oregon Department of Fish and Wildlife, Portland, OR.

ODFW. 2005. Oregon threatened and endangered species list [online]. Wildlife Division, Oregon Department of Fish and Wildlife, Salem, OR. Updated 9/1/2005. [Cited 2006.] Available from: http://www.dfw.state.or.us/wildlife/diversity/threatened_endangered.asp.

Ondreicka R, Ginter E, Kortus J. 1966. Chronic toxicity of aluminium in rats and mice and its effects on phosphorus metabolism. *Brit J Industr Med* 23:305-312.

Orcutt HG. 1950. The life history of the starry flounder *Platichthys stellatus* (Pallas). Fish bulletin no. 78. Bureau of Marine Fisheries, Division of Fish and Game, California Department of Natural Resources, Sacramento, CA.

Oregon Natural Heritage Information Center. 2004. Rare, threatened and endangered species of Oregon. Oregon Natural Heritage Information Center, Institute for Natural Resources, Oregon State University, Portland, OR.

Oregonian. 2006. China's mercury flushes into Oregon's rivers. The Oregonian, Portland, OR, November 24, 2006.

Page LM, Burr BM. 1991. A field guide to freshwater fishes of North America north of Mexico. Peterson Field Guides, Houghton Mifflin Company, Boston, MA.

Parsons J. 2001. Aquatic plant sampling protocols. Pub. no. 01-03-017. Environmental Assessment Program, Washington Department of Ecology, Olympia, WA.

Pastorok RA, Bartell SM, Ferson S, Ginzburg LR, eds. 2001. Ecological modeling in risk assessment: chemical effects on populations, ecosystems, and landscapes. Lewis Publishers, Boca Raton, FL.

Patton JF, Dieter MP. 1980. Effects of petroleum hydrocarbons on hepatic function in the duck. *Comp Biochem Physiol* 65C:33-36.

Peakall DB. 1974. Effects of di-n-butylphthalate and di-2-ethylhexylphthalate on the eggs of ring doves. *Bull Environ Contam Toxicol* 12:698-702.

Pennak R. 1978. Fresh water invertebrates of the U.S. 2nd ed. John Wiley & Sons Inc., New York, NY.

Pennington T. 2002. Personal communication on July 7, 2002 (telephone conversation with Maryann Welsch, Windward Environmental LLC, regarding aquatic vegetation in the ISA). ANS Research Assistant, Center for Lakes and Reservoirs, Portland State University, Portland, OR.

Phillips DJH, Rainbow PS. 1989. Strategies of trace metal sequestration in aquatic organisms. *Mar Environ Res* 28(207-210).

Pierce EP. 1993. The breeding biology and behavior of the purple sandpiper (*Calidris maritima*) in Svalbard. Master's thesis. University of Bergen, Bergen, Norway.

Pinkney AE, Harshbarger JC, May EB, Melancon MJ. 2000. Tumor prevalence and biomarkers in brown bullheads (*Ameiurus nebulosus*) from the tidal Potomac River watershed. CBFO-C99-04. US Fish and Wildlife Service, Annapolis, MD.

Pinkney AE, Harshbarger JC, May EB, Melancon MJ. 2004a. Tumor prevalence and biomarkers of exposure and response in brown bullhead (*Ameiurus nebulosus*) from the Anacostia River, Washington, DC and Tuckahoe River, Maryland, USA. *Environ Toxicol Chem* 23(3):638-347.

Pinkney AE, Harshbarger JC, May EB, Melancon MJ. 2004b. Tumor prevalence and biomarkers of exposure in brown bullhead (*Ameiurus nebulosus*) from Back River, Furnace Creek, and Tuckahoe River, Maryland. *Arch Environ Contam Toxicol* 46:492-501.

Plant Conservation Alliance. 2009. Purple loosestrife, *Lythrum salicaria* L. Plant Conservation Alliance's Alien Plant Working Group least wanted [online]. National Park Service, Washington, DC. Updated 7/7/09. [Cited 2/7/11.] Available from: <http://www.nps.gov/plants/alien/fact/lysa1.htm>.

Poole AF. 1983. Courtship feeding, clutch size, and egg size in ospreys: a preliminary report. In: Bird DM, Seymour NR, Gerrard JM, eds, Biology and management of bald eagles and ospreys. Harpell Press, Saint-Anne-de-Bellevue, Quebec, pp 243-256.

Precht E, Huettel M. 2003. Rapid wave-driven advective pore water exchange in a permeable coastal sediment. J Sea Res 51:93-107.

Puchy CA, Marshall DB. 1993. Oregon wildlife diversity plan 1993-1998. Oregon Department of Fish and Wildlife, Portland, OR.

Racey GD, Euler DL. 1983. Changes in mink habitat and food selection as influenced by cottage development in central Ontario. J Appl Ecol 20:387-402.

Raimondo S, Montague B, Barron MG. 2007. Determinants of variability in acute to chronic toxicity ratios for aquatic invertebrates and fish. Environ Toxicol Chem 26(9):2019-2023.

Raleigh RF, Hickman T, Solomon RC, Nelson PC. 1984. Habitat suitability information: Rainbow trout. FWS/OBS-82/10.60. US Fish and Wildlife Service, Washington, DC.

Rasmussen AD, Banta GT, Andersen O. 2000. Cadmium dynamics in estuarine sediments: effects of salinity and lugworm bioturbation. Environ Toxicol Chem 19(2):380-386.

Reed PB. 1996. National list of plant species that occur in wetlands: 1996 national summary [online]. US Fish and Wildlife Service, Washington, DC. Updated 1/24/97. [Cited 2003.] Available from: <http://www.nwi.fws.gov/bha/download/1996/national.txt>.

Rehnberg BC, Schreck CB. 1986. Acute metal toxicology of olfaction in coho salmon: behavior receptors, and odor-metal complexation. Bull Environ Contam Toxicol 36:579-86.

Restum JC, Bursian SJ, Giesy JP, Render JA, Helferich WG, Shipp EB, Verbrugge DA. 1998. Multigenerational study of the effects of consumption of PCB-contaminated carp from Saginaw Bay, Lake Huron, on mink. 1. Effects on mink reproduction, kit growth and survival, and selected biological parameters. J Toxicol Environ Health 54(A):343-375.

Rhoads DC, Germano JD. 1986. Interpreting long-term changes in benthic community structure: a new protocol. Hydrobiologia 142:291-308.

Rice CA, Myers MS, Willis ML, French BL, Casillas E. 2000. From sediment bioassay to fish biomarker – connecting the dots using simple trophic relationships. Mar Environ Res 50:527-533.

Riisgard HU, Larsen PS. 2005. Water pumping and analysis of flow in burrowing zoobenthos: an overview. Aqua Ecol 39:237-258.

Roberts MH, Hargis WJ, Strobel CJ, DeLisle PF. 1989. Acute toxicology of PAH contaminated sediments to the estuarine fish, *Leiostomus xanthurus*. Bull Environ Contam Toxicol 42:142-149.

Roling JA, Bain LJ, Gardea-Torresdey J, Bader J, Baldwin WS. 2006. Hexavalent chromium reduces larval growth and alters gene expression in mummichog (*Fundulus heteroclitus*). *Environ Toxicol Chem* 25(10):2725-2733.

RSET. 2009. Regional sediment evaluation framework for the Pacific Northwest. Final. May 2009. Regional Sediment Evaluation Team (US Army Corps of Engineers - Seattle District, Portland District, Walla Walla District, and Northwestern Division; US Environmental Protection Agency Region 10; Washington Department of Ecology; Washington Department of Natural Resources; Oregon Department of Environmental Quality; Idaho Department of Environmental Quality; National Marine Fisheries Service; US Fish and Wildlife Service).

Saban L, Andersen H. 2004. Memorandum to Eric Blischke and Chip Humphrey, US Environmental Protection Agency Region 10, on behalf of Lower Willamette Group, regarding shorebird beach sampling. July 16, 2004. Windward Environmental LLC, Seattle, WA.

Sample BE, Suter GW, II. 1999. Ecological risk assessment in a large river-reservoir: 4. Piscivorous wildlife. *Environ Toxicol Chem* 18(4):610-620.

Sample BE, Opreko DM, Suter GW. 1996. Toxicological benchmarks for wildlife. 1996 revision. ES/ERM-86/R3. Office of Environmental Management, US Department of Energy, Washington, DC.

Sandahl JF, Baldwin DH, Jenkins JJ, Scholz NL. 2004. Odor-evoked field potentials as indicators of sublethal neurotoxicity in juvenile coho salmon (*Oncorhynchus kisutch*) exposed to copper, chlorpyrifos, or esfenvalerate. *Can J Fish Aquat Sci* 61:404-413.

Sandahl JF, Miyasaka G, Koide N, Ueda H. 2006. Olfactory inhibition and recovery in chum salmon (*Oncorhynchus keta*) following copper exposure. *Can J Fish Aquat Sci* 63:1840-1847.

Sandahl JF, Baldwin DH, Jenkins JJ, Scholz NL. 2007. A sensory system at the interface between urban stormwater runoff and salmon survival. *Environ Sci Technol* 41:2998-3004.

Sandercock FK. 1991. Life history of coho salmon (*Oncorhynchus kisutch*). In: Groot C, Margolis L, eds, Pacific salmon life histories. UBC Press, Vancouver, BC, pp 395-46.

Sanders HO, Mayer FL, Jr, Walsh DF. 1973. Toxicity, residue dynamics, and reproductive effects of phthalate esters in aquatic invertebrates. *Environ Res* 6:84-90.

Sanderson JT, Van den Berg M. 1999. Toxic equivalency factors (TEFs) and their use in ecological risk assessment: a successful method when used appropriately. *Human Ecol Risk Assess* 5:43-52.

Santore RC, Di Toro DM, Paquin PR, Allen HE, Meyer JS. 2001. Biotic ligand model of the acute toxicity of metals. 2. Application to acute copper toxicity in freshwater fish and *Daphnia*. *Environ Toxicol Chem* 20(10):2397-2402.

Satoh H, Nakamura Y, Okabe S. 2007. Influences of infaunal burrows on the community structure and activity of ammonia-oxidizing bacteria in intertidal sediments. *App Environ Microbiol* 73(4):1341-1348.

Saucier D, Astic L, Rioux P. 1991. The effects of early chronic exposure to sublethal copper on the olfactory discrimination of rainbow trout, (*Oncorhynchus mykiss*). *Environ Biol Fish* 30:345-351.

Saunders RL, Sprague JB. 1967. Effects of copper-zinc mining pollution on spawning migration of Atlantic salmon. *Wat Res* 1:419-432.

Schimmel SC, Patrick Jr JM, Forester J. 1977. Toxicity and bioconcentration of BHC and lindane in selected estuarine animals. *Arch Environ Contam Toxicol* 6:355-363.

Schlatterer B, Coenen TMM, Ebert E, Grau R, Hilbig V, Munk R. 1993. Effects of bis(tri-n-butyltin) oxide in Japanese quail exposed during egg laying period: an interlaboratory comparison study. *Arch Environ Contam Toxicol* 24:440-448.

Schmitt CJ, Dethloff GM, eds. 2000. Biomonitoring of environmental status and trends (BEST) program: selected methods for monitoring chemical contaminants and their effects in aquatic ecosystems. USGS/BRD-2000-0005. Biological Resources Division, US Geological Survey, Columbia, MO.

Schreck CB, Snelling JC, Ewing RE, Bradford CS, Davis LE, Slater CH. 1994a. Migratory behavior of adult spring chinook salmon in the Willamette River and its tributaries. Completion report. Prepared for Bonneville Power Administration, Portland, OR. Oregon Cooperative Fishery Research Unit, Oregon State University, Corvallis, OR.

Schreck CB, Snelling JC, Ewing RE, Bradford CS, Davis LE, Slater CH. 1994b. Migratory behavior of juvenile spring chinook salmon in the Willamette River. Completion report. Prepared for Bonneville Power Administration, Portland, OR. Oregon Cooperative Fishery Research Unit, Oregon State University, Corvallis, OR.

Scott WB, Crossman EJ. 1973. Freshwater fishes of Canada. Bulletin 184. Fisheries Research Board of Canada, Ottawa, ON, Canada.

SEA, Windward, Anchor, Kennedy/Jenks. 2002. Portland Harbor remedial investigation/feasibility study round 1 field sampling plan. Draft. Prepared for Lower Willamette Group. Striplin Environmental Associates, Inc., Olympia, WA; Windward Environmental LLC, Seattle, WA; Anchor Environmental, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR.

SEA. 2002. Portland Harbor remedial investigation/feasibility study sediment profile image survey of the Lower Willamette River. Prepared for Lower Willamette Group. Striplin Environmental Associates, Inc., Olympia, WA.

SEA, Fishman, Ellis, Windward, Anchor, Kennedy/Jenks. 2003. Portland Harbor remedial investigation/feasibility study round 1 field sampling report. Prepared for Lower Willamette Group. Striplin Environmental Associates, Inc., Olympia, WA; Fishman Environmental Services, Portland, OR; Ellis Environmental Services, Estacada, OR; Windward Environmental LLC, Seattle, WA; Anchor Environmental, Seattle, WA; Kennedy/Jenks Consultants, Portland, OR.

Sedell JR, Froggatt JL. 1984. Importance of streamside forests to large rivers: The isolation of the Willamette River, Oregon, USA, from its floodplain by snagging and streamside forest removal. *Verh Internat Verein Limnol* 22:1828-1834.

Shephard B. 2008. Copper effects on fish behavior: a critical review and synopsis of existing studies. Draft white paper. Office of Environmental Assessment, US Environmental Protection Agency Region 10, Seattle, WA.

Shephard B, Zodrow J. 2009. Copper effects on fish behavior: a review and synopsis of existing studies. PowerPoint presentation at 30th SETAC North America Annual Meeting, New Orleans, LA, November 23, 2009. US Environmental Protection Agency Region 10, Seattle, WA.

Shephard B. 2010. Copper effects on fish behavior: a review and synopsis of existing studies. PowerPoint presentation at International Copper Association Decennial Science Symposium, Paradise Valley, AZ, July 23, 2010. US Environmental Protection Agency Region 10, Seattle, WA.

Smith SL, MacDonald DD, Keenleyside KA, Ingersoll CG, Field LJ. 1996. A preliminary evaluation of sediment quality assessment values for freshwater ecosystems. *J Great Lakes Res* 22:624-638.

Soster FM, McCall PL. 1990. Benthos response to disturbance in western Lake Erie: field experiments. *Can J Fish Aquat Sci* 47:1970-1985.

Spalding MG, Frederick PC, McGill HC, Bouton SN, McDowell LR. 2000. Methylmercury accumulation in tissues and its effects on growth and appetite in captive great egrets. *J Wildl Dis* 36(3):411-422.

Sparling DW. 2000. Ecotoxicology of organic contaminants to amphibians. In: Sparling DW, Linder G, Bishop C, eds, *Ecotoxicology of amphibians and reptiles*. SETAC, Pensacola, FL, pp 461-495.

Sparling DW, Linder G, Bishop CA, eds. 2000. *Ecotoxicology of amphibians and reptiles*. SETAC Technical Publications Series, Ingersoll CG, ed. Society of Environmental Toxicology and Chemistry (SETAC) Press, Pensacola, FL.

Spehar RL. 1976. Cadmium and zinc toxicity to flagfish, *Jordanella floridae*. *J Fish Res Bd Can* 33:1939-1945.

- Sprague JB. 1964. Avoidance of copper-zinc solutions by young salmon in the laboratory. *J Water Pollut Control Fed* 36:990-1004.
- SREL. 1999. Terrestrial toxicity reference values (TRVs). Manual ERD-AG-03. 04/06/99. Environmental Restoration Division, Savannah River Ecology Laboratory, Aiken, SC.
- Stalmaster MV, Gessaman JA. 1984. Ecological energetics and foraging behavior of overwintering bald eagles. *Ecol Monogr* 54:407-428.
- Stephan CE, Mount DI, Hansen DJ, Gentile JH, Chapman GA, Brungs WA. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. PB85-227049. Office of Research and Development, US Environmental Protection Agency, Washington, DC.
- Stern JH, Hennessy DP, Patmont CR. 2003. Improving estimates of contaminant exposure for mobile organisms: an assessment of area-weighted home range exposure estimates applied to the relationship between sediment chemistry and liver lesions in English sole. Puget Sound Research Conference 2003, Vancouver, BC.
- Stier DJ, Crance JH. 1985. Habitat suitability index models and instream flow suitability curves: American shad. *FWS Biol Rep* 82(10.88). US Fish and Wildlife Service, Washington, DC.
- Stuber RJ, Gebhart G, Maughan OE. 1982. Habitat suitability index models: Bluegill. *FWS/OBS-82/10.8* [online]. US Fish and Wildlife Service, Washington, DC. [Cited 3/28/02.] Available from: <http://www.nwrc.gov/wdb/pub/hsi/hsi-008.pdf>.
- Suter GW, Tsao CL. 1996. Toxicological benchmarks for screening potential contaminants of concern for effects on aquatic biota: 1996 revision. Prepared for U.S. Department of Energy Office of Environmental Management. Risk Assessment Program, Health Sciences Research Division.
- Tacon AGJ. 1992. Nutritional fish pathology: morphological signs of nutrient deficiency and toxicity in farmed fish. *FAO fish technical paper no. T330*. Fisheries and Aquaculture Department, Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.
- Tetra Tech. 1993. Willamette River basin water quality study. Willamette River ecological systems investigation component report. Submitted to Oregon Department of Environmental Quality, Portland, OR. Tetra Tech, Inc., Redmond, WA.
- Tetra Tech. 1995. Willamette River basin water quality study. Phase II. Ecological monitoring component: assessment of aquatic communities and biological indices. Submitted to Oregon Department of Environmental Quality, Portland, OR. Tetra Tech, Inc., Redmond, WA.
- Thorp JP, Covich AP. 2001. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, NY.

Tillitt DE, Gale RW, Meadows JC, Zajicek JL, Peterman PH, Heaton SN, Jones PD, Bursian SJ, Kubiak TJ, Giesy JP, Aulerich RJ. 1996. Dietary exposure of mink to carp from Saginaw Bay. 3. Characterization of dietary exposure to planar halogenated hydrocarbons, dioxin equivalents, and biomagnification. *Environ Sci Technol* 30:283-291.

Trotter PC. 1997. Sea-run cutthroat trout: life history profile. In: Sea-run cutthroat trout: biology, management, and future conservation. Oregon Chapter, American Fisheries Society, Portland, OR, pp 7-15.

Turner JL. 1966. Distribution and food habits of centrarchid fishes in the Sacramento-San Joaquin delta. In: Turner JL, Kelly DW, eds, Fish Bulletin 136. California Department of Fish and Game, Sacramento, CA, pp 144-153.

USDA FS. 1990. *Fraxinus latifolia* Benth., Oregon ash. In: Silvics of North America: 2. Hardwoods. Agriculture Handbook 654. US Department of Agriculture, Forest Service, Washington, DC. http://www.na.fs.fed.us/pubs/silvics_manual/volume_2/fraxinus/latifolia.htm.

USDA FS. 2011. Fire Effects Information System database: *Iris pseudacorus*, *Typha latifolia*, *Cornus sericea*, *Spiraea douglasii*, *Salix lucida* subsp. *lasianдра* [online]. Rocky Mountain Research Station, USDA Forest Service, Missoula, MT. [Cited 2/7/11.] Available from: <http://www.fs.fed.us/database/feis>.

USEPA. 1993. Wildlife exposure factors handbook. EPA/600/R-93/187a. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

USFWS. 1961. Sturgeons. Fishery leaflet 526. Bureau of Commercial Fisheries, US Fish and Wildlife Service, Washington, DC.

USFWS. 2002. Recovery plan for the California red-legged frog (*Rana aurora draytonii*). US Fish and Wildlife Service Region 1, Portland, OR.

USFWS. 2007. Procedure for sampling fish, collecting tissues, and conducting an external fish health assessment. Oregon Fish and Wildlife Office, US Fish and Wildlife Service, Portland, OR.

USFWS. 2011. Species profile: Water howellia (*Howellia aquatilis*) [online]. Environmental Conservation Online System (ECOS), US Fish and Wildlife Service, Washington, DC. [Cited 2/7/11.] Available from: <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?scode=Q2RM>.

USGS. 2002. Illustrated field guide for assessing external and internal anomalies in fish. Information and technology report USGS/BRD/ITR-2002-0007 [online]. US Geological Survey, Washington, DC. Available from: http://www.cerc.usgs.gov/pubs/center/pdfDocs/ITR_2002_0007.pdf.

USGS. 2004. Personal communication on unspecified date (R. Grove to Jeremy Buck, US Fish and Wildlife Service, Portland, OR) data on otter body weights in western Oregon and Washington). US Geological Survey, Portland, OR.

USGS. 2006. NASQAN (National Stream Quality Accounting Network) Program. Statistical summaries of published data: Columbia River Basin [online]. US Geological Survey, Portland, OR. Updated 4/13/2006. [Cited 1/2/07.] Available from:
<http://water.usgs.gov/nasqan/data/statsumtxt.html>.

Van den Berg M, Birnbaum L, Bosveld ATC, Brunström B, Cook P, Feeley M, Giesy JP, Hanberg A, Hasegawa R, Kennedy S, Kubiak T, Larsen JC, van Leeuwen FXR, Djen Liem AK, Nolt C, Peterson RE, Poellinger L, Safe S, Schrenk D, Tillitt D, Tysklind M, Younes M, Waern F, Zacharewski T. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ Health Perspect* 106(12):775-792.

Veinott G, Northcote T, Rosenau M, Evan RD. 1999. Concentrations of strontium in the pectoral fin rays of the white sturgeon (*Acipenser transmontanus*) by laser ablation sampling- inductively coupled plasma - mass spectrometry as an indicator of marine migrations. *Can J Fish Aquat Sci* 56:1981-1990.

Vile JS, Friesen TA. 2005. Description and categorization of nearshore habitat in the Lower Willamette River. In: Friesen TA, ed, Biology, behavior, and resources of resident and anadromous fish in the Lower Willamette River. Final report of research, 2000-2004. Prepared for City of Portland. Oregon Department of Fish and Wildlife, Clackamas, OR, pp 17-62.

Vogel S. 1994. Life in moving fluids: the physical biology of flow. Second ed. Princeton University Press, Princeton, NJ.

Vogelbein WK, Unger M. 2003. The Elizabeth River monitoring program 2001-2002: Association between mummichog liver histopathology and sediment chemical contamination. Prepared for Virginia Department of Environmental Quality. Virginia Institute of Marine Science, The College of William and Mary, Gloucester Point, VA.

Wang W, Gorsuch JW, Hughes JS, eds. 1997. Plants for environmental studies. CRC Press LLC, Lewis Publishers, Boca Raton, FL.

Ward DA, Farr RA. 1989. Effects of waterway development on anadromous and resident fish in Portland Harbor. Annual progress report (1988). Fish Research Project, Oregon. Oregon Department of Fish and Wildlife, Portland, OR.

Ward DA, Farr RA. 1990. Effects of waterway development on anadromous and resident fish in Portland Harbor. Annual progress report (1989). Fish Research Project, Oregon. Oregon Department of Fish and Wildlife, Portland, OR.

Ward DA, Knutsen CJ. 1991. Effects of waterway development on anadromous and resident fish in Portland Harbor. Annual progress report (1990). Fish Research Project, Oregon. Oregon Department of Fish and Wildlife, Portland, OR.

Ward DA, Connolly PJ, Farr RA, Nigro AA. 1988. Feasibility of evaluating the impacts of waterway development on anadromous and resident fish in Portland Harbor. Annual progress report, Fish Research Project, Oregon. Oregon Department of Fish and Wildlife, Portland.

Ward DA, Nigro AA, Farr RA, Knutsen CJ. 1994. Influence of waterway development on migrational characteristics of juvenile salmonids in the lower Willamette River, Oregon. N Am J Fish Manage 14:362-371.

Ward DL, Nigro AA. 1992. Differences in fish assemblages among habitats found in the lower Willamette River, Oregon: application of and problems with multivariate analysis. Fish Res 13:119-132.

Ward DL, Nigro AA, Farr RA, Knutsen CJ. 1992. Categorization of waterway developments in the lower Willamette River, Oregon, based on risks to juvenile salmonids. Oregon Department of Fish and Wildlife, Clackamas, OR.

Ward DP, Smal CM, Fairley JS. 1986. The food of mink (*Mustela vison*) in the Irish midlands. Proc Royal Irish Acad 86b:169-182.

Warnock SE, Takekawa JY. 1995. Habitat preferences of wintering shorebirds in a temporally changing environment: western sandpipers in the San Francisco Bay estuary. Auk 112(4):920-930.

WDFW. 2007. Letter dated March 8, 2007 to T. Do, Windward, from M. Wall, WDFW, regarding information on a tagged white sturgeon. March 8, 2007. Columbia River Sturgeon Project, Washington Department of Fish and Wildlife Region 5, Vancouver, WA.

WDG. 1980. Study of impacts of project modification and river regulation on riparian habitats and associated wildlife along the Columbia River. Washington Department of Game, Olympia, WA.

Weber JB, Mrozek E, Jr. 1979. Polychlorinated biphenyls: Phytotoxicity, absorption and translocation by plants, and inactivation by activated carbon. Bull Environ Contam Toxicol 23:412-17.

Wells KD. 2007. The ecology and behavior of amphibians. University of Chicago Press, Chicago, IL.

Widdows J, Page DS. 1993. Effects of tributyltin and dibutyltin on the physiological energetics of the mussel, *Mytilus edulis*. Mar Environ Res 35:233-249.

Wiemeyer SN. 1991. Personal communication on (estimate of average adult bald eagle body weights) cited in EPA's Wildlife Exposures Handbook (1993).

Wiemeyer SN, Lamont TG, Bunck CM, Sindelar CR, Gramlich F, Fraser JD, Byrd MA. 1984. Organochlorine pesticide, polychlorobiphenyl, and mercury residues in bald eagle eggs-1969-1979-and their relationships to shell thinning and reproduction. Arch Environ Contam Toxicol 13:529-549.

Wiemeyer SN, Bunck CM, Stafford CJ. 1993. Environmental contaminants in bald eagle eggs 1980-84 and further interpretations of relationships to productivity and shell thickness. Arch Environ Contam Toxicol 24:213-227.

Windward. 2004. Portland Harbor Superfund Site ecological risk assessment: comprehensive synopsis of approaches and methods. Draft. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005a. Portland Harbor RI/FS. Ecological preliminary risk evaluation. WE-05-0007. Draft. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA.

Windward. 2005b. Portland Harbor RI/FS. Round 2 sampling of invertebrates using multiplate samplers. Field sampling report. WE-05-0010. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006a. Portland Harbor RI/FS Round 3 sampling for lamprey (*Lampetra* sp.) tissue field sampling report. Prepared for the Lower Willamette Group. Windward Environmental LLC, Seattle, WA.

Windward. 2006b. Portland Harbor Superfund Site: Proposed ecological risk assessment decision framework. Draft. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA.

Windward. 2007. Lower Duwamish Waterway remedial investigation. Baseline ecological risk assessment. Prepared for Lower Duwamish Waterway Group. Windward Environmental LLC, Seattle, WA.

Windward, Integral. 2008. Portland Harbor RI/FS. Round 3 lamprey ammocoete phase 2 toxicity testing data report. Draft. WE-08-0003. Prepared for Lower Willamette Group. May 7, 2008. Windward Environmental LLC, Seattle, WA.

Windward. 2009a. Portland Harbor RI, Appendix G: Baseline ecological risk assessment. Draft. Prepared for the Lower Willamette Group. August 19, 2009. Windward Environmental LLC, Seattle, WA.

Windward. 2009b. Portland Harbor RI/FS bioaccumulation modeling report. Draft. WW09-0003. Prepared for Lower Willamette Group. July 21, 2009. Windward Environmental LLC, Seattle, WA.

Windward, Integral. 2005a. Portland Harbor RI/FS. Ecological risk assessment technical memorandum: results of field sampling reconnaissance for invertebrates using a benthic sledge, bongo net, diaphragm pump, and Schindler trap. WE-05-0005. Draft. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA; Integral Consulting, Inc., Mercer Island, WA.

Windward, Integral. 2005b. Portland Harbor RI/FS. Field sampling plan: Round 2 sampling of benthic invertebrate tissue. WE-05-0008. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA; Integral Consulting, Inc., Mercer Island, WA.

Windward, Avocet, TerraStat. 2006. Portland Harbor Superfund Site ecological risk assessment. Interpretive report: estimating risks to benthic organisms using predictive models based on sediment toxicity tests. Draft. WE-06-0002. Prepared for Lower Willamette Group. Windward Environmental LLC, Seattle, WA; Avocet Consulting, Kenmore, WA; TerraStat Consulting Group, Snohomish, WA.

Winsor MH, Boese BL, Lee H, Randall RC, Specht DT. 1990. Determination of the ventilation rates of interstitial and overlying water by the clam *Macoma nasuta*. Environ Toxicol Chem 9:209-213.

Wisconsin DNR. 2004. Species information: Common cattail (*Typha latifolia*) [online]. Wisconsin Department of Natural Resources, Madison, WI. Updated 9/2/04. [Cited 2/7/11.] Available from: http://dnr.wi.gov/invasives/fact/com_cattail.htm.

Wise MH, Linn IJ, Kennedy CR. 1981. A comparison of the feeding biology of mink *Mustela vison* and otter *Lutra lutra*. J Zool London 195:181-213.

WNHP. 1999. Field guide to selected rare plants of Washington: *Lomatium bradshawii* (Bradshaw's lomatium) [online]. Washington Natural Heritage Program, Olympia, WA. [Cited 2/5/11.] Available from: <http://www1.dnr.wa.gov/nhp/refdesk/fguide/pdf/lobr.pdf>.

Wren CD, Hunter DB, Leatherland JF, Stokes PM. 1987. The effects of polychlorinated biphenyls and methylmercury, singly and in combination on mink. II: Reproduction and kit development. Arch Environ Contam Toxicol 16:449-454.

Wydoski RS, Whitney RR. 1979. Inland fishes of Washington. University of Washington Press, Seattle, WA.

Wydoski RS, Whitney RR. 2003. Inland fishes of Washington. 2nd ed. University of Washington Press, Seattle, WA.

Zhou JM, Qin F, Cong L, Xu B. 2004. Toxicity of PCBs (Aroclor-1221, 1254) to embryos and larvae of *Xenopus laevis*. Bull Environ Contam Toxicol 73:379-384.